

COMPLEXITY OF ATRIODIC CONTINUA

by

CHRISTOPHER TODD KENNAUGH, M.S.

A DISSERTATION

IN

MATHEMATICS

Submitted to the Graduate Faculty
of Texas Tech University in
Partial Fulfillment of
the Requirements for
the Degree of

DOCTOR OF PHILOSOPHY

APPROVED BY

Wayne Lewis (Chair)

Robert Byerly

Razvan Gelca

Fred Hartmeister,

Dean of the Graduate School

May, 2009

©2009, Christopher Kennaugh

ACKNOWLEDGMENTS

Thanks to Wayne Lewis for being the author's teacher and advisor and for directing this dissertation. Thanks to the Department of Mathematics and Statistics at Texas Tech for having the author as a graduate student and teaching assistant. Thanks to Robert Byerly and Razvan Gelca for being on the committee of this dissertation. Thanks to Dale Daniel for getting the author in to topology. Thanks to the author's folks Chris and Pat for all their support. Dedicated to the Golden Triangle, Texas.

CONTENTS

ACKNOWLEDGMENTS	ii
ABSTRACT	iv
LIST OF TABLES	v
LIST OF FIGURES	vi
I. BASIC TERMINOLOGY	1
II. INTRODUCTION	3
III. PRELIMINARIES	7
IV. THE EXAMPLE $_nK$	12
4.1 Definitions	13
4.2 The Bonding Map $_n\phi$	17
4.3 Factoring	36
V. FURTHER QUESTIONS	68
BIBLIOGRAPHY	70
APPENDIX: SOME BONDING MAPS	73

ABSTRACT

This dissertation investigates the relative complexity between a continuum and its proper subcontinua (see [19]), in particular, providing examples of atriodic n -od-like continua. Let X be a continuum and n be an integer greater than or equal to three. If X is homeomorphic to an inverse limit of simple- n -od graphs with simplicial bonding maps and is simple- $(n - 1)$ -od-like, it is shown that the bonding maps can be simplicially factored through a simple- $(n - 1)$ -od. This implies, in particular, that X is homeomorphic to an inverse limit of simple- $(n - 1)$ -od graphs with simplicial bonding maps. This factoring is subsequently used (in a strategy adapted from [15]) to show that a specific inverse limit of simple- n -ods with simplicial bonding maps, having the property of every proper nondegenerate subcontinuum being an arc, is not simple- $(n - 1)$ -od-like.

LIST OF TABLES

4.1	Definition of ${}_nX_0$	17
4.2	Definition of ${}_nX_1$	17
4.3	Definition of ${}_n\phi$	18
4.4	Subgraphs of ${}_nX_1$ corresponding to vertices of $D({}_n\phi, {}_nX_1)$	25
4.5	Determination of $D({}_n\phi, {}_nX_1)$	25
4.6	Determination of $d[{}_n\phi]$	26
4.7	Edge inverses under $d[{}_n\phi]$	27
4.8	Definition of ${}_nY_1$	27
4.9	Definition of ${}_n\lambda$	28

LIST OF FIGURES

3.1	A possibility for $\mathcal{W}$ and $\mathcal{V}$	8
3.2	A possibility for $\mathcal{U}$	9
4.1	The bonding map ${}_3\phi$	19
4.2	The bonding map ${}_4\phi$	20
4.3	The bonding map ${}_5\phi$	21
4.4	The map $d[{}_5\phi]$	26
4.5	Factoring of ${}_5\Phi_0^m$	63
4.6	Factoring of ${}_5\Gamma_1^m$	67
A.1	Ingram bonding map	73
A.2	Davis-Ingram bonding map	74
A.3	Minc bonding map	75

CHAPTER I

BASIC TERMINOLOGY

Definition 1.1. A *continuum* is a connected, compact, metric space.

Definition 1.2. A continuum is *decomposable* if it is the union of two of its proper subcontinua and is *indecomposable* otherwise.

Definition 1.3. A continuum is *hereditarily decomposable* (*hereditarily indecomposable*) if each of its nondegenerate subcontinua is decomposable (indecomposable).

Definition 1.4. A continuum X is a *triad* (3-od) if it contains a subcontinuum M so that $X \setminus M$ is the union of three nonempty mutually separated sets.

Definition 1.5. A continuum is *atriodic* if it does not contain a triad.

Definition 1.6. For a positive integer n , a *simple- n -od* is the union of n arcs joined at an end point.

Definition 1.7. For a sequence of (*factor*) spaces $\{X_i\}_{i=0}^{\infty}$ and sequence of (*bonding*) maps $\{f_i\}_{i=0}^{\infty}$ where $f_i : X_{i+1} \rightarrow X_i$, the *inverse limit* is the subspace $M = \{(x_0, x_1, \dots) \in \prod_{i=0}^{\infty} X_i : f_i(x_{i+1}) = x_i \text{ for all } i\}$, denoted $M = \varprojlim \{X_i, f_i^j\}_{i=0, j \geq i}^{\infty}$, where $f_i^i = \text{identity on } X_i$ and for $j > i$ $f_i^j = f_i \circ f_{i+1} \circ \dots \circ f_{j-1}$.

Proposition 1.8. Let X be a continuum and $\mathcal{P}$ be a collection of connected, compact polyhedra. Then the following are equivalent:

- 1) $X \approx \varprojlim \{X_i, f_i^j\}_{i=0, j \geq i}^{\infty}$ where $X_i \in \mathcal{P}$ and f_i is a continuous surjection for each i .
- 2) For each $\epsilon > 0$ there exist $Y \in \mathcal{P}$ and a continuous surjection $f : X \rightarrow Y$ so that $\text{diam}(f^{-1}(y)) < \epsilon$ for each $y \in Y$.
- 3) For each $\epsilon > 0$ there exists an open covering $\mathcal{U}$ of X whose nerve is a member of $\mathcal{P}$ and so that $\text{diam}(U) < \epsilon$ for each $U \in \mathcal{U}$.

Definition 1.9. A continuum satisfying the conditions of the above proposition is $\mathcal{P}$ -like.

Definition 1.10. A continuum which is arc-like (simple-1-od-like, simple-2-od-like) is *chainable* (also *snake-like*).

Definition 1.11. A continuum is *subchainable* if every proper subcontinuum is chainable.

Definition 1.12. Let n be a positive integer greater than or equal to 3, K be a continuum, and $v \in K$. Then v is a *branch point of K of order n* if and only if for each $\epsilon > 0$ there exists an open cover $\mathcal{U}$ of K so that $\text{mesh}(\mathcal{U}) < \epsilon$, the nerve of $\mathcal{U}$ is a simple- n -od, and v is in the element of $\mathcal{U}$ of order n .

Definition 1.13. A *graph* is a one-dimensional, connected, finite simplicial complex. If G is a graph, then $V(G)$ denotes the set of vertices and $E(G)$ denotes the set of edges of G .

Definition 1.14. A map f between graphs is *simplicial* provided each edge is either mapped linearly onto an edge or mapped into a single vertex.

Definition 1.15 (Def. 2.1 [9]). A graph H is *simpler* ($\leq$) than a graph G if there exists a simplicial monotone map from G onto H .

Definition 1.16. For graphs G and H , a G -like continuum X , and an H -like continuum Y , Y is *simpler* than X if H is simpler than G .

CHAPTER II

INTRODUCTION

In 1951, R.H. Bing showed in [3] that among hereditarily decomposable tree-like continua, chainability is equivalent to atriodicity and stated the following question in [2] (M denotes the pseudo-arc): “It would be interesting to know if each nondegenerate bounded hereditarily indecomposable plane continuum which does not separate the plane is homeomorphic to M . This question would be answered in the affirmative if it were shown that each bounded atriodic plane continuum which does not separate the plane can be chained.” In the same year, the following statement, along with other related claims, appeared in an abstract [1] given by R.D. Anderson: “The author demonstrates the existence of a hereditarily indecomposable plane continuum, not separating the plane, which is not homeomorphic to a pseudo-arc (that is, a chained hereditarily indecomposable plane continuum).” The first published example of such a continuum, in 1979, is contained in [8] by W.T. Ingram, constructed with a modification to the first published counterexample to Bing’s second question, given in [7], with crookedness appropriately built-in so as to retain nonchainability.

Also by Ingram and published in 1972, [7] is an example of an inverse limit of simple-3-ods with a single bonding map, where nonchainability and atriodicity are implied by the example’s properties of positive span and of every proper nondegenerate subcontinuum being an arc, respectively. A. Lelek, in the 1964 publication [10], defined for a metric space X the *span* $\sigma(X)$ as the least upper bound on the numbers ϵ for which there exists a connected subspace $Z \subseteq X \times X$ with $\pi_1(Z) = \pi_2(Z)$ so that the distance between a and b is greater than or equal to ϵ for each $(a, b) \in Z$ (*surjective span* $\sigma^*(X)$ has the above definition with the additional condition that $\pi_1(Z) = X$) and showed that if a continuum X is chainable then $\sigma(X) = 0$ (the converse is presently open). More generally, for a space X , metric space Y , and a map f of X into Y , the *span of f* $\sigma(f)$ is the least upper bound on the numbers ϵ for which there exists a connected subspace $Z \subseteq X \times X$ with $\pi_1(Z) = \pi_2(Z)$ so that

the distance between $f(a)$ and $f(b)$ is greater than or equal to ϵ for each $(a, b) \in Z$. Then, $\sigma(X)$ is the span of the identity map on X . Ingram showed that for the bonding maps f_i^j in the example from [7], there exists a positive number ϵ so that $\sigma(f_0^j) > \epsilon$ for each j and that this implies positive span of the inverse limit. In the 1968 publication [6], Ingram demonstrated that the property of being atriodic, known to be present in chainable continua, also holds, more generally, among subchainable continua. The nature of the single bonding map in [7] ensures that any nondegenerate subcontinuum of the inverse limit which is not an arc is not proper, establishing atriodicity of the inverse limit. Having every proper nondegenerate subcontinuum as an arc also implies, by the aforementioned result of Bing, that the Ingram example [7], being nonchainable, is necessarily indecomposable. Other inverse limits with the previously discussed properties in common with [7] have been constructed utilizing similar techniques as in [7] in demonstrating positive span, such as in the example by J.F. Davis and Ingram [5]. Published in 1988, this continuum also has the property of admitting a (monotone) map to a chainable continuum with only one nondegenerate point inverse which is an arc.

Noting that a simple- n -od-like continuum is simple- $(n + 1)$ -od-like, for example, some properties of a continuum can be more readily realized when constructing that continuum as a more “complex” space. Such instances are in [11] by W. Lewis and in [13] by J.C. Mayer, published in 1983. In [11], for each n , an inverse limit of simple- n -ods is constructed by Lewis with bonding maps sufficiently varying so as to produce chainability of the inverse limit and whose symmetry allows for a homeomorphism on the inverse limit to be induced, having only one fixed point and every other point with period n . Thereby, construction of a continuum as one which is simple- n -od-like facilitates a description of a period n homeomorphism on that continuum, controlled so as to be simpler than would initially appear in the construction, being simple-2-od-like. Thus is given such a homeomorphism for each n on a chainable continuum and, further, by introducing crookedness, such a homeomorphism for each n on the pseudo-arc. In [13], an inverse limit of simple-4-ods, referred to as the “X-odic” continuum,

is constructed by Mayer with a single bonding map, having the previously discussed properties of [7] and utilizing similar techniques in demonstrating positive span as [7], in addition to admitting an embedding in the plane with a Lake-of-Wada channel. In the same publication as [13], in [18] S.W. Young showed that the bonding map in [13] can be factored through a simple-triod, giving that the simple-4-od-like X-odic continuum is simple-3-od-like. Young remarks, “although the bonding map,” induced from the factoring, used in the representation of the (X-odic) continuum X in the simpler form “does not seem to help in establishing the main properties of the continuum X , it must inevitably detract from its name.”

Also by Young and in the same publication as [13] and [18], [19] has the following beginning to its introduction: “One of the remarkable features of the continuum of W.T. Ingram ([7]) is the ‘gap’ in complexity between the continuum and its proper subcontinua. Specifically, the continuum is T-like (simple-triod-like), not arc-like and every proper subcontinuum is arc-like. This combination of structural properties leads us to ask if there is a continuum with an even wider ‘gap’.” Question 1 follows: “Does there exist a continuum which is 4-od-like, not simple-triod-like and every proper subcontinuum is arc-like?” Similar questions with slight variations on this question are Problem 115 from [12] published in 1983, “Is there a continuum which is 4-od-like, not T-like, and every nondegenerate proper subcontinuum of which is an arc?” and Problem 5 from [4] published in 1990, “Does there exist an atriodic simple-4-od-like continuum which is not simple-triod-like?,” with a positive answer to the second being a positive answer to all three.

These questions were answered in the affirmative by P. Minc with the example in [15] published in 1993. In response to these questions, Minc states in [15], “Even after a perfunctory glance at the problems, it becomes apparent that they should have a positive answer. It is very easy to get an example of a simple-4-od-like continuum such that every proper subcontinuum is an arc. Most of such continua appear not to be simple-triod-like and it is very likely that they really are not. So the only difficulty is a proof,” where “a topological invariant different than the span is needed

to distinguish between those continua which are simple-triod-like and those that are not. Another way of approaching the problem is to use a continuum with simplicial bonding maps and prove that they cannot be factored through a simple triod.” The following is the abstract from [14] by Minc and published in 1994: “An operation d on simplicial maps between graphs is introduced and used to characterize simplicial maps which can be factored through an arc. The characterization yields a new technique of showing that some continua are not chainable,” demonstrated with the examples in [7] and [5], “and allows to prove that span zero is equivalent to chainability for inverse limits of trees with simplicial bonding maps.” Theorem 3.3 from [14] establishes that the following are equivalent for an inverse limit X of trees with simplicial bonding maps f_i^j :

- 1) X is chainable
- 2) $\sigma^*(X) = 0$
- 3) for each i there exists $j > i$ so that f_i^j can be (simplicially) factored through an arc.

This alternate, combinatorial technique involving factoring and the operation d is extended by Minc to the example in [15] in demonstrating that it is not simple-3-od-like, where an inverse limit of simple-4-ods is constructed with a single bonding map, controlled so that every proper nondegenerate subcontinuum is an arc and so that the bonding map f_0^j cannot be (simplicially) factored through a simple-3-od for each j .

The purpose of this dissertation is to investigate this gap in the complexity between a continuum and its proper subcontinua further, in particular when the proper nondegenerate subcontinua are arcs. For each integer $n \geq 3$, an inverse limit ${}_nK$ of simple- n -ods with a single bonding map ${}_n\phi$ is constructed which is not simple- $(n-1)$ -od-like and whose every proper nondegenerate subcontinuum is an arc, established in Chapter IV. To this end, an extension and generalization of the strategy employed in [15] is developed with a general theorem on factoring established in Chapter III.

CHAPTER III

PRELIMINARIES

In this chapter, a theorem on factoring is given (Theorem 3.3) stating that for certain inverse limits in the simplicial setting, in the case the inverse limit is simpler (Definition 1.16) than the complexity of the factor spaces would indicate, the bonding maps are able to be factored through simpler graphs. In light of Lemma 3.1, the proof of Theorem 3.3 follows from the proof of Proposition 2.1 in [15].

Let K be a continuum and n be an integer greater than or equal to 3.

Lemma 3.1. *Suppose $\mathcal{V}$ is an open cover of K so that the nerve of $\mathcal{V}$ is a simple- n -od and $\mathcal{W}$ is an open cover of K refining $\mathcal{V}$ so that the nerve of $\mathcal{W}$ is a simple- $(n-1)$ -od. Then, there exists an amalgamation $\mathcal{U}$ of $\mathcal{W}$ refining $\mathcal{V}$ so that the nerve of $\mathcal{U}$ is a simple- $\tilde{n}$ -od for some $\tilde{n} \in \{1, \dots, n-1\}$ and U_0 , the element of $\mathcal{U}$ of order $\tilde{n}$, is contained in V_0 , the element of $\mathcal{V}$ of order n .*

Proof. For each $i \in \{1, \dots, n\}$ and some $\nu_i \in \{1, 2, \dots\}$, let $V_0^i, V_1^i, V_2^i, \dots, V_{\nu_i}^i$ be a linear chain in $\mathcal{V}$ so that $\mathcal{V} = \{V_j^i : i \in \{1, \dots, n\} \text{ and } j \in \{0, \dots, \nu_i\}\}$, where $V_0^i = V_0$ for each $i \in \{1, \dots, n\}$, and let $V^i = \bigcup_{j=1}^{\nu_i} V_j^i$. For each $i \in \{1, \dots, n-1\}$ and some $\mu_i \in \{1, 2, \dots\}$, let $W_0, W_1^i, W_2^i, \dots, W_{\mu_i}^i$ be a linear chain in $\mathcal{W}$ so that $\mathcal{W} = \{W_0, W_j^i : i \in \{1, \dots, n-1\} \text{ and } j \in \{1, \dots, \mu_i\}\}$, where W_0 is the element of $\mathcal{W}$ of order $n-1$.

Suppose $W_0 \not\subseteq V_0$, since otherwise the claim holds with $\mathcal{U} = \mathcal{W}$. Let $\tilde{i} \in \{1, \dots, n\}$ and $\tilde{j} \in \{1, \dots, \nu_{\tilde{i}}\}$ so that $W_0 \subseteq V_{\tilde{j}}^{\tilde{i}}$ and $W_0 \not\subseteq V_{\tilde{j}-1}^{\tilde{i}}$. For each $i \in \{1, \dots, n-1\}$, let $j_i \in \{1, \dots, \mu_i\}$ so that $\bigcup_{j=1}^{j_i} W_j^i \subseteq V^{\tilde{i}} \cup V_0$ and $W_{j_i+1}^i \not\subseteq V^{\tilde{i}} \cup V_0$ or $W_{j_i+1}^i$ does not exist. Let $\tilde{j}_m = \max\{j \in \{0, \dots, \nu_{\tilde{i}}\} : W_k^i \subseteq V_j^{\tilde{i}} \text{ and } W_k^i \not\subseteq V_{j-1}^{\tilde{i}} \text{ if } V_{j-1}^{\tilde{i}} \text{ exists or } W_k^i \not\subseteq V^{\tilde{i}} \text{ for all } i \neq \tilde{i} \text{ if } V_{j-1}^{\tilde{i}} \text{ does not exist, for some } i \in \{1, \dots, n-1\} \text{ and } j \in \{1, \dots, j_i\}\}$, and let $i' \in \{1, \dots, n-1\}$ and $k_{i'} \in \{1, \dots, j_{i'}\}$ so that $W_{k_{i'}}^{i'} \subseteq V_{\tilde{j}_m}^{\tilde{i}}$ and $W_{k_{i'}}^{i'} \not\subseteq V_{\tilde{j}_m-1}^{\tilde{i}}$ if $V_{\tilde{j}_m-1}^{\tilde{i}}$ exists or $W_{k_{i'}}^{i'} \not\subseteq V^{\tilde{i}}$ for all $i \neq \tilde{i}$ if $V_{\tilde{j}_m-1}^{\tilde{i}}$ does not exist.

Without loss of generality, suppose

i) there exists $\hat{i} \in \{1, \dots, n-1\}$ so that $\{i \in \{1, \dots, n-1\} \setminus \{i'\} : W_{j_i+1}^i \text{ does not exist}\} = \{\hat{i}, \hat{i}+1, \dots, n-1\}$ or

ii) $\{i \in \{1, \dots, n-1\} \setminus \{i'\} : W_{j_i+1}^i \text{ does not exist}\} = \emptyset$.

Let $\tilde{n} = \hat{i} - 1$ if (i) and $\tilde{n} = n - 1$ if (ii). Let $\tilde{j}_M = \tilde{j}$ if $\tilde{j} \geq \tilde{j}_m$, or $\tilde{j}_M = \tilde{j}_m - 1$ if $\tilde{j} < \tilde{j}_m$ and $\{W_k^i \subseteq V_{\tilde{j}_m}^i \text{ and } W_k^i \not\subseteq V_{\tilde{j}_m-1}^i : i \in \{1, \dots, n-1\} \setminus \{i'\} \text{ and } k \in \{1, \dots, j_i\}\} \cup \{W_k^{i'} \subseteq V_{\tilde{j}_m}^i \text{ and } W_k^{i'} \not\subseteq V_{\tilde{j}_m-1}^i : k \in \{1, \dots, k_{i'}-1\}\} = \emptyset$, or $\tilde{j}_M = \tilde{j}_m$ if otherwise.

For each $i \in \{1, \dots, \tilde{n}\} \setminus \{i'\}$, let $U_1^i, U_2^i, \dots, U_{\mu_i-j_i}^i$ be a linear chain of open sets defined as $U_j^i = W_{j_i+j}^i$ for each $j \in \{1, \dots, \mu_i-j_i\}$, and let U_0 be an open set defined as $U_0 = \bigcup(\{W_j^i \subseteq V_0 : i \in \{1, \dots, n-1\} \setminus \{i'\} \text{ and } j \in \{1, \dots, j_i\}\} \cup \{W_j^{i'} \subseteq V_0 : j \in \{1, \dots, k_{i'}-1\}\})$, and let $U_1^{i'}, U_2^{i'}, \dots, U_{\tilde{j}_M+\mu_{i'}-k_{i'}+1}^{i'}$ be a linear chain of open sets defined as $U_j^{i'} = \bigcup(\{W \subseteq V_j^i \text{ and } W \not\subseteq V_{j-1}^i : W \in \{W_0, W_k^i : i \in \{1, \dots, n-1\} \setminus \{i'\} \text{ and } k \in \{1, \dots, j_i\}\}\} \cup \{W_k^{i'} \subseteq V_j^i \text{ and } W_k^{i'} \not\subseteq V_{j-1}^i : k \in \{1, \dots, k_{i'}-1\}\})$ for each $j \in \{1, \dots, \tilde{j}_M\}$ and $U_{\tilde{j}_M+j}^{i'} = W_{k_{i'}-1+j}^{i'}$ for each $j \in \{1, \dots, \mu_{i'}-k_{i'}+1\}$. Then the claim holds with $\mathcal{U} = \{U_0\} \cup \{U_j^i : i \in \{1, \dots, \tilde{n}\} \setminus \{i'\} \text{ and } j \in \{1, \dots, \mu_i-j_i\}\} \cup \{U_j^{i'} : j \in \{1, \dots, \tilde{j}_M+\mu_{i'}-k_{i'}+1\}\}$. $\square$





Figure 3.1: A possibility for $\mathcal{W}$ and $\mathcal{V}$.


 Figure 3.2: A possibility for $\mathcal{U}$.

Proposition 3.2. *Suppose $v \in K$ is a branch point of K of order n and K is simple- $(n-1)$ -od-like. Then, for each $\epsilon > 0$, there exists an open cover $\mathcal{U}$ of K so that $\text{mesh}(\mathcal{U}) < \epsilon$, the nerve of $\mathcal{U}$ is a simple- $\tilde{n}$ -od for some $\tilde{n} \in \{1, \dots, n-1\}$, and v is within ϵ of the element of $\mathcal{U}$ of order $\tilde{n}$.*

Proof. The claim follows by applying Lemma 3.1 and Definition 1.12 □

Theorem 3.3. *Suppose K is homeomorphic to $\varprojlim \{X_i, \Phi_i^j\}_{i=0, j \geq i}^\infty$ where X_i is a simple- n -od graph and Φ_i is simplicial for each $i \in \{0, 1, \dots\}$. If K is simple- $(n-1)$ -od-like, then for each $i \in \{0, 1, \dots\}$ there exist $j \in \{i+1, i+2, \dots\}$, a simple- $\tilde{n}$ -od T for some $\tilde{n} \in \{1, \dots, n-1\}$, and simplicial maps $\alpha : X_j \rightarrow T$ and $\beta : T \rightarrow X_i$ so that $\beta \circ \alpha = \Phi_i^j$ and $\beta(t_0) = x_0$ where t_0 and x_0 are the branch points of T and X_i , respectively.*

Proof. Let $\{B_x : x \in V(X_i)\}$ be a collection of mutually exclusive connected open sets in X_i so that $x \in B_x$ for each $x \in V(X_i)$, and so the nerve of $\mathcal{V} = \{\pi_i^{-1}(V) : V \in$

$\{B_x : x \in V(X_i)\} \cup \{\text{open edges of } X_i\}$ is a simple- n -od. Since K is simple- $(n-1)$ -od-like, by Lemma 3.1 there exists an open cover $\mathcal{U}$ of K refining $\mathcal{V}$ so that the nerve of $\mathcal{U}$ is a simple- $\tilde{n}$ -od for some $\tilde{n} \in \{1, \dots, n-1\}$ and U_0 , the element of $\mathcal{U}$ of order $\tilde{n}$, is contained in $\pi_i^{-1}(B_{x_0})$.

Let $\mathcal{U}_B = \{U \in \mathcal{U} : U \subseteq \pi_i^{-1}(B_x) \text{ for some } x \in V(X_i)\}$ and for each $U \in \mathcal{U}_B$ let $x_U \in V(X_i)$ so that $U \subseteq \pi_i^{-1}(B_{x_U})$. Define an equivalence relation $\equiv$ on $\mathcal{U}_B$ as $U_p \equiv U_q$ if and only if $x_U = x_{U_p}$ for all $U \in \langle U_p, U_q \rangle \cap \mathcal{U}_B$, where $\langle U_p, U_q \rangle$ denotes the unique linear chain in $\mathcal{U}$ between U_p and U_q , for each $U_p, U_q \in \mathcal{U}_B$. Let T be the graph defined as $V(T) = \mathcal{U}_B / \equiv$ and $\widehat{U_p}, \widehat{U_q} \in V(T)$ are adjacent if and only if $U \in \widehat{U_p}$ or $U \in \widehat{U_q}$ for each $U \in \langle U_p, U_q \rangle \cap \mathcal{U}_B$ for some $U_p \in \widehat{U_p}$ and $U_q \in \widehat{U_q}$. Since $U_0 \subseteq \pi_i^{-1}(B_{x_0})$, $U_0 \in \mathcal{U}_B$ and $x_{U_0} = x_0$. Let $\widehat{U_0} \in V(T)$ so that $U_0 \in \widehat{U_0}$. Then, if $\widehat{U_p}, \widehat{U_q} \in V(T)$ are adjacent or equal and $U_0 \in \langle U_p, U_q \rangle$ for some $U_p \in \widehat{U_p}$ and $U_q \in \widehat{U_q}$, then $\widehat{U_p} = \widehat{U_0}$ or $\widehat{U_q} = \widehat{U_0}$. Thus, $\widehat{U_0}$ is the unique vertex of T with order greater than 2, provided T is not an arc, and without loss of generality has order $\tilde{n}$. Then, T is a simple- $\tilde{n}$ -od.

Define the piecewise-linear map $\beta : T \rightarrow X_i$ as $\beta(t) = x_U$ for some $U \in t$ for each $t \in V(T)$. Suppose $t_p, t_q \in V(T)$ are adjacent, and let $U_p \in t_p$ and $U_q \in t_q$. If x_{U_p} and x_{U_q} are not adjacent and not equal then there exists $x \in V(X_i) \setminus \{x_{U_p}, x_{U_q}\}$ so that $x \in \langle x_{U_p}, x_{U_q} \rangle$, where $\langle x_{U_p}, x_{U_q} \rangle$ denotes the unique arc in X_i between x_{U_p} and x_{U_q} , and so $x_U = x$ for some $U \in \langle U_p, U_q \rangle \cap \mathcal{U}_B$, contradicting $U \in t_p \cup t_q$. Thus, $\beta(t_p)$ and $\beta(t_q)$ are adjacent or equal, and so β is simplicial. Also, $\beta(t_0) = x_{U_0} = x_0$ where $t_0 = \widehat{U_0}$.

Let ϵ be a Lebesgue number for $\mathcal{U}$ and $j > i$ so that the diameter of $\pi_j^{-1}(x)$ is less than ϵ for each $x \in X_j$. For each $x \in V(X_j)$ let $U_x \in \mathcal{U}$ so that $\pi_j^{-1}(x) \subseteq U_x$. Since $U_x \cap \pi_i^{-1}(\Phi_i^j(x)) \neq \emptyset$, $U_x \subseteq \pi_i^{-1}(B_{\Phi_i^j(x)})$ for each $x \in V(X_j)$, and so $U_{\tilde{x}} \in \mathcal{U}_B$ with $x_{U_{\tilde{x}}} = \Phi_i^j(\tilde{x})$ for each $\tilde{x} \in V(X_j)$. Let $t_{\tilde{x}} \in V(T)$ so that $U_{\tilde{x}} \in t_{\tilde{x}}$, and define the piecewise-linear map $\alpha : X_j \rightarrow T$ as $\alpha(x) = t_{\tilde{x}}$ for each $x \in V(X_j)$. Then, $\beta \circ \alpha(\tilde{x}) = \beta(t_{\tilde{x}}) = x_{U_{\tilde{x}}} = \Phi_i^j(\tilde{x})$ for all $\tilde{x} \in V(X_j)$. Suppose $x_p, x_q \in V(X_j)$ are adjacent and t_{x_p} and t_{x_q} are not adjacent and not equal. Then there exist $t \in V(T) \setminus \{t_{x_p}, t_{x_q}\}$

so that $t \in \langle t_{x_p}, t_{x_q} \rangle$ and $U_1 \in \langle U_{x_p}, U_{x_q} \rangle$ so that $U_1 \in t$. Let $W_1, W_2, \dots, W_m \subseteq \langle x_p, x_q \rangle$ for some $m \in \{1, 2, \dots\}$ be a linear chain of connected sets covering $\langle x_p, x_q \rangle$ with $x_p \in W_1$ so that the diameter of $\pi_j^{-1}(W_k)$ is less than ϵ for each $k \in \{1, \dots, m\}$. Since $U \cap \pi_j^{-1}(W_k) \neq \emptyset$ for some $k \in \{1, \dots, m\}$ for all $U \in \langle U_{x_p}, U_{x_q} \rangle$, $U \cap \pi_i^{-1}(\langle \Phi_i^j(x_p), \Phi_i^j(x_q) \rangle) \neq \emptyset$ for all $U \in \langle U_{x_p}, U_{x_q} \rangle$, and so if $U \in \langle U_{x_p}, U_{x_q} \rangle \cap \mathcal{U}_B$ then $U \subseteq \pi_i^{-1}(B_{\Phi_i^j(x_p)})$ or $U \subseteq \pi_i^{-1}(B_{\Phi_i^j(x_q)})$. If $\Phi_i^j(x_p) = \Phi_i^j(x_q)$ then $t_{x_p} = t_{x_q}$, giving a contradiction. Thus, without loss of generality, $\Phi_i^j(x_p) \neq \Phi_i^j(x_q)$, $x_{U_1} = \Phi_i^j(x_p)$, and there exists $U_2 \in \langle U_{x_p}, U_1 \rangle \cap \mathcal{U}_B$ so that $x_{U_2} = \Phi_i^j(x_q)$. Let $k \in \{1, \dots, m\}$ so that $U_2 \cap \pi_j^{-1}(W_k) \neq \emptyset$. Since $\pi_i(U_2) \subseteq B_{\Phi_i^j(x_q)}$, $\Phi_i^j(W_k) \cap B_{\Phi_i^j(x_q)} \neq \emptyset$, and so by linearity $\pi_i^{-1}(W) \cap B_{\Phi_i^j(x_q)} \neq \emptyset$ for all $W \in \langle W_k, W_m \rangle$. But, $\pi_j^{-1}(W) \subseteq U_1$ for some $W \in \langle W_k, W_m \rangle$, and so $U_1 \cap \pi_i^{-1}(B_{\Phi_i^j(x_q)}) \neq \emptyset$ contradicting $U_1 \subseteq \pi_i^{-1}(B_{\Phi_i^j(x_p)})$. Thus, $\alpha(x_p)$ and $\alpha(x_q)$ are adjacent or equal, and so α is simplicial. $\square$

Corollary 3.4. *Suppose K is homeomorphic to $\varprojlim \{X_i, \Phi_i^j\}_{i=0, j \geq i}^\infty$ where X_i is a simple- n -od graph and Φ_i is simplicial for each $i \in \{0, 1, \dots\}$. If K is simple- $(n-1)$ -od-like, then K is homeomorphic to $\varprojlim \{Y_i, \Psi_i^j\}_{i=0, j \geq i}^\infty$ where Y_i is a simple- $\tilde{n}$ -od graph, for some $\tilde{n} \in \{1, \dots, n-1\}$, and Ψ_i is simplicial for each $i \in \{0, 1, \dots\}$.*

Proof. By Theorem 3.3, there exist a strictly increasing sequence $\{n_0, n_1, \dots\} \subseteq \{0, 1, \dots\}$ with $n_0 = 0$ and for each $i \in \{0, 1, \dots\}$ simplicial maps $\alpha_i : X_{n_{i+1}} \longrightarrow Y_i$ and $\beta_i : Y_i \longrightarrow X_{n_i}$ with Y_i a simple- $\tilde{n}$ -od for some $\tilde{n} \in \{1, \dots, n-1\}$ so that $\beta_i \circ \alpha_i = \Phi_{n_i}^{n_{i+1}}$. For each $i \in \{0, 1, \dots\}$ define $\Psi_i : Y_{i+1} \longrightarrow Y_i$ as $\Psi_i = \alpha_i \circ \beta_{i+1}$, and the claim holds. $\square$

CHAPTER IV

THE EXAMPLE ${}_nK$

In this chapter, for an integer n greater than or equal to 3, an example ${}_nK$ of an inverse limit of simple- n -ods with a single bonding map ${}_n\phi$ is given. As in [7], [5], and [15], characteristics of the bonding map ensure every proper nondegenerate subcontinuum being an arc, implying atriodicity of the continuum. The bonding map for ${}_nK$ being a simplicial map from a graph which is a subdivision of the range graph, as for the other examples, allows for the inverse limit to be represented “simplicially” where the factor spaces are graphs with fixed sets of vertices and the bonding maps are simplicial. In this context, as developed by Minc in [14], certain combinatorial conditions involving the operation d and factoring allow for the determination of the nonchainability of the inverse limit. If the bonding map defining the inverse limit is particularly “nice”, then these conditions are satisfied, implying that the bonding maps cannot be factored through arcs, implying nonchainability.

Critical to this program with factoring bonding maps through arcs (simple-2-ods), when considering the dual of the factoring, the dual of the arc remains an arc (or is a point or is empty). When factoring through a simple-3-od, for example, this is not necessarily the case, where the dual of the simple-3-od may be more “complicated” than a simple-3-od. To maintain this control of complexity when considering an extension of Minc’s algorithm with such factorings, particulars of the specific bonding maps in question need to be invoked. Such is done by Minc in [15], where, again in the simplicial setting, a combinatorial program involving the factoring of bonding maps through simple-3-ods is adapted.

For the specific example in [15], the nature of the defining bonding map allows for a certain factoring to be arranged, provided the existence of a factoring in general, thereby avoiding a “bad” case where the dual of the simple-3-od could possibly be more “complicated.” Specifically, branch point of the simple-3-od going to branch point can be avoided in the factoring, implying, for the dual of simple-3-ods, there

being at most one point of order bigger than 2. In extending Minc's program further, a strategy to control complexity is also used in the factoring of the bonding maps of ${}_nK$ through simple- $(n - 1)$ -ods. In addition to satisfying Minc's combinatorial conditions, the nature of the defining bonding map ${}_n\phi$ induces characteristics in the bonding maps that allow for the rearrangement of the factoring, given by the theorem in the previous chapter, to one in which the dual of the simple- $(n - 1)$ -od remains a simple- $(n - 1)$ -od. This is accomplished in conjunction with and facilitated by branch point of the simple- $(n - 1)$ -od going to branch point in the factoring as given by the theorem, which, paradoxically, is the worst-case scenario with regard to allowing for the greatest possible complexity of the dual of the simple- $(n - 1)$ -od.

4.1 Definitions

Definition 4.1 (Def. 2.1 [14]). For a graph G , let $D(G)$ denote the graph such that

- i) $V(D(G))$ and $E(G)$ are in one-to-one correspondence and
- ii) two vertices of $D(G)$ are adjacent if and only if the edges of G corresponding to the vertices have a non-empty intersection.

For $v \in V(D(G))$, v^* denotes the edge of G corresponding to v .

Definition 4.2 (Def. 2.4 [14]). Let $\phi : G_1 \longrightarrow G_0$ be a simplicial map between graphs, $C = \{c \subseteq G_1 : c \text{ is a component of } \phi^{-1}(e) \text{ and } \phi(c) = e \text{ for some } e \in E(G_0)\}$, and $D(\phi, G_1)$ be the graph so that

- i) $V(D(\phi, G_1))$ and C are in one-to-one correspondence and
- ii) two vertices of $D(\phi, G_1)$ are adjacent if and only if the subgraphs of G_1 corresponding to the vertices have a non-empty intersection.

For $v \in V(D(\phi, G_1))$, v^* denotes the subgraph of G_1 corresponding to v .

Let $d[\phi] : D(\phi, G_1) \longrightarrow D(G_0)$ be the simplicial map determined by $d[\phi](v) = w$ where $w \in V(D(G_0))$ so that $\phi(v^*) = w^*$, for every $v \in V(D(\phi, G_1))$.

Definition 4.3 (Def. 2.10 [14]). Let $\phi : G_1 \longrightarrow G_0$ and $\psi : G_2 \longrightarrow G_1$ be simplicial maps between graphs. Let $d[\phi, \psi] : D(\phi \circ \psi, G_2) \longrightarrow D(\phi, G_1)$ be the simplicial map

determined by $d[\phi, \psi](v) = w$ where w is the vertex of $D(\phi, G_1)$ so that $\psi(v^*) \subseteq w^*$, for every vertex v of $D(\phi \circ \psi, G_2)$.

Definition 4.4 (Def. 4.1 [14]). Let $\phi : G_1 \longrightarrow G_0$ be a simplicial map between graphs. Then ϕ is *ultra light* if it is light and v^* is an edge of G_1 for each $v \in V(D(\phi, G_1))$.

Definition 4.5 (Def. 5.1 [14]). A graph G' *subdivides* a graph G if $V(G) \subseteq V(G')$ and for each $e \in E(G)$ there is an arc $(e, G') \subseteq G'$ so that

- i) (e, G') has the same endpoints as e ,
- ii) $(d, G') \cap (e, G') = d \cap e$ for $d, e \in E(G)$ and $d \neq e$, and
- iii) for each $v \in V(G')$, $v \in (e, G')$ for some $e \in E(G)$, and for each $e' \in E(G')$, e' is an edge of (e, G') for some $e \in E(G)$.

If $v \in V(G)$ and $e \in E(G)$ so that $v \in e$, then (v, e, G') denotes the edge of (e, G') containing v .

Proposition 4.1 (Prop. 5.2 [14]). *If G' is a graph subdividing a graph G and G'' is a graph subdividing G' , then G'' subdivides G .*

Definition 4.6 (Def. 5.3 [14]). Let G'_0 and G'_1 be graphs subdividing the graphs G_0 and G_1 , respectively, and $\phi : G_1 \longrightarrow G_0$ and $\phi' : G'_1 \longrightarrow G'_0$ be simplicial maps so that $\phi'(v) = \phi(v)$ for all $v \in V(G_1)$. If for all $e \in E(G_1)$, (e, G'_1) is an edge whenever $\phi(e)$ is degenerate, and ϕ' restricted to (e, G'_1) is an isomorphism onto $(\phi(e), G'_0)$ whenever $\phi(e)$ is nondegenerate, then ϕ' is a *subdivision of ϕ matching G'_0* .

Proposition 4.2 (Prop. 5.4 [14]). *Let $\phi : G_1 \longrightarrow G_0$ be a simplicial map between graphs. Let G'_0 be a graph subdividing G_0 . Then there is a subdivision ϕ' of ϕ matching G'_0 , unique up to an isomorphism.*

Definition 4.7 (Def. 5.5 [14]). Let G be a graph and $S : V(G) \longrightarrow \{X \subseteq E(G) : X \neq \emptyset\}$. Then S is an *edge selection on G* if for all $v \in V(G)$, $v \in e$ for each $e \in S(v)$.

Definition 4.8 (Def. 5.5 [14]). Let G_0 and G_1 be graphs, S be an edge selection on G_1 , G'_1 be a graph subdividing G_1 , and $\phi : G'_1 \rightarrow G_0$ be simplicial. If there exist a graph H_1 and a simplicial map $\lambda : H_1 \rightarrow D(\phi, G'_1)$ so that H_1 is a subdivision of G_1 and λ is an isomorphism with

- i) $(v, e, G'_1) \subseteq [\lambda(v)]^*$ for each $v \in V(G_1)$ and each $e \in S(v)$ and
- ii) $[\lambda(v)]^* \subseteq (e, G'_1)$ for each $e \in E(G_1)$ and $v \in V((e, H_1)) \setminus V(G_1)$,

then ϕ is consistent on S and λ is a consistency isomorphism.

Definition 4.9 (Def. 5.7 [14]). Let G_0 and G_1 be graphs with edge selections S_0 and S_1 , respectively, G'_1 be a graph subdividing G_1 , and $\phi : G'_1 \rightarrow G_0$ be simplicial. Then, ϕ preserves (S_0, S_1) provided that

- i) $\phi((v, e, G'_1)) \in S_0(\phi(v))$ for each $v \in V(G_1)$ and each $e \in S_1(v)$ and
- ii) for each $e, e' \in E(G'_1)$ so that $e \neq e'$ and $e \cap e' = v$ for some $v \in V(G'_1)$, either $\phi(e) \in S_0(\phi(v))$ or $\phi(e') \in S_0(\phi(v))$.

Definition 4.10 (Def. 5.10 [14]). Let $\{G_i\}_{i=0}^\infty$ and $\{G'_i\}_{i=1}^\infty$ be collections of graphs and $\Phi = \{\phi_i\}_{i=0}^\infty$ be a collection of simplicial maps so that for each $i \in \{1, 2, \dots\}$, G'_i is a subdivision of G_i and $\phi_{i-1} : G'_i \rightarrow G_{i-1}$. By Proposition 4.2 there exists a collection of simplicial maps $\{\psi_i\}_{i=0}^\infty$ so that $\psi_0 = \phi_0$ and for each $i \in \{1, 2, \dots\}$, ψ_i is a subdivision of ϕ_i matching the domain of ψ_{i-1} . Let $X_0 = G_0$, for each $i \in \{1, 2, \dots\}$, X_i be the domain of ψ_{i-1} , and for $i, j \in \{0, 1, \dots\}$ so that $j > i$ let $\Phi_i^j = \psi_i \circ \psi_{i+1} \circ \dots \circ \psi_{j-1}$ and $\Phi_i^i = \text{identity on } X_i$. Then we say $\varprojlim \{X_i, \Phi_i^j\}_{i=0, j \geq i}^\infty$ is generated by the sequence Φ .

Let $S = \{S_i\}_{i=1}^\infty$ where for each $i \in \{1, 2, \dots\}$, S_i is an edge selection on G_i . We say Φ preserves S if ϕ_i preserves (S_i, S_{i+1}) for each $i \in \{1, 2, \dots\}$.

Definition 4.11 (Def. 5.10 [14]). We say two inverse limits $\varprojlim \{X_i, \Phi_i^j\}_{i=0, j \geq i}^\infty$ and $\varprojlim \{Y_i, \Psi_i^j\}_{i=0, j \geq i}^\infty$ with simplicial bonding maps are isomorphic if there exists a sequence of isomorphisms $\{\lambda_i\}_{i=0}^\infty$ so that $\lambda_i : X_i \rightarrow Y_i$ and $\lambda_i \circ \Phi_i^j = \Psi_i^j \circ \lambda_j$ for all $i, j \in \{0, 1, \dots\}$ with $j \geq i$.

Definition 4.12. Let $x_0, x_1, \dots, x_{m_x}$ for some $m_x \in \{1, 2, \dots\}$ be vertices of some graph G and define $\langle x_0, x_1, \dots, x_{m_x} \rangle$ to be the path in G given by $x_0, x_1, \dots, x_{m_x}$, respecting order, if and only if x_i and x_{i+1} are adjacent for all $i \in \{0, \dots, m_x - 1\}$. Let $y_0, y_1, \dots, y_{m_y}$ for some $m_y \in \{1, 2, \dots\}$ be vertices of G so that y_i and y_{i+1} are adjacent for all $i \in \{0, \dots, m_y - 1\}$, and define $\langle x_0, x_1, \dots, x_{m_x} \rangle \vee \langle y_0, y_1, \dots, y_{m_y} \rangle = \langle x_0, x_1, \dots, x_{m_x}, y_1, \dots, y_{m_y} \rangle$, if and only if $x_{m_x} = y_0$.

4.2 The Bonding Map ${}_n\phi$

In this section, a single bonding map ${}_n\phi$ between simple- n -ods is defined for an inverse limit ${}_nK$. Characteristics of the map ensure that every proper nondegenerate subcontinuum of ${}_nK$ is an arc (Proposition 4.3) and satisfy certain conditions essential in the implementation of the previously mentioned strategy of Minc. In particular, ${}_n\phi$ is well-behaved in the sense of inducing no folding in $d[{}_n\phi]$ (Proposition 4.7) and with respect to satisfying conditions sufficient for demonstrating a “subdivision” of ${}_nK$ as being isomorphic with the dual of ${}_nK$ (Propositions 4.6 and 4.9). Also, a certain symmetry is exhibited by the bonding maps (Proposition 4.12), used in the next section to obtain a “well-behaved” factoring.

For what follows, let ${}_nX_0$ be a simple- n -od and ${}_nA_0^1, {}_nA_0^2, \dots, {}_nA_0^n$ be n arcs so that $\bigcup_{i=1}^n {}_nA_0^i = {}_nX_0$ and ${}_nA_0^i \cap {}_nA_0^j = \{v_0\}$ for each distinct $i, j \in \{1, \dots, n\}$, and so v_0 is an endpoint of ${}_nA_0^i$ for each $i \in \{1, \dots, n\}$.

Let ${}_nX_0$ be the graph defined in the following table.

arc	as a subgraph
${}_nA_0^i$	$\langle v_0, v_i \rangle$ for each $i \in \{1, \dots, n-2\}$
${}_nA_0^{n-1}$	$\langle v_0, v_{n-1}, v_{n+1} \rangle$
${}_nA_0^n$	$\langle v_0, v_n, v_{n+2} \rangle$

Table 4.1: Definition of ${}_nX_0$

Let ${}_nX_1$ be a subdivision of ${}_nX_0$, where ${}_nA_1^i$ is a subdivision of ${}_nA_0^i$ for each $i \in \{1, \dots, n\}$ defined in the following table.

arc	as a subgraph
${}_nA_1^i$	$\langle v_0, u_i, u_{(n-3)+i}, u_{2(n-3)+i}, \dots, u_{2(n-1)(n-3)+i}, v_i \rangle$ for each $i \in \{1, \dots, n-3\}$
${}_nA_1^{n-2}$	$\langle v_0, u_{(2n-1)(n-3)+2(n-1)+1}, v_{n-2} \rangle$
${}_nA_1^{n-1}$	$\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2}, u_{(2n-1)(n-3)+2(n-1)+3}, v_{n-1}, v_{n+1} \rangle$
${}_nA_1^n$	$\langle v_0, u_{(2n-1)(n-3)+1}, \dots, u_{(2n-1)(n-3)+2(n-1)}, v_n, v_{n+2} \rangle$

Table 4.2: Definition of ${}_nX_1$

Let ${}_n\phi : {}_nX_1 \longrightarrow {}_nX_0$ be the light simplicial map determined by the following table where ${}_n\phi(v)$ is defined for $v \in V({}_nX_1)$.

v	${}_n\phi(v)$
v_0	v_0
v_i	v_{n+2} for all $i \in \{1, \dots, n-3\}$
v_{n-2}	v_{n+1}
v_{n-1}	v_n
v_n	v_n
v_{n+1}	v_{n+2}
v_{n+2}	v_{n+2}
u_i	v_{n-1} for all $i \in \{1, \dots, n-3\}$
$u_{j(n-3)+i}$	v_0 for all $i \in \{1, \dots, n-3\}$ and for all odd $j \in \{1, \dots, 2(n-1)-1\}$
$u_{j(n-3)+i}$	$v_{i-1+\frac{j}{2} \bmod n-2}$ for all $i \in \{1, \dots, n-3\}$ and for all even $j \in \{2, \dots, 2(n-1)-2\}$
$u_{2(n-1)(n-3)+i}$	v_n for all $i \in \{1, \dots, n-3\}$
$u_{(2n-1)(n-3)+1}$	v_{n-1}
$u_{(2n-1)(n-3)+i}$	v_0 for all even $i \in \{2, \dots, 2(n-1)\}$
$u_{(2n-1)(n-3)+3}$	v_{n-2}
$u_{(2n-1)(n-3)+i}$	$v_{\frac{i-3}{2}}$ for all odd $i \in \{5, \dots, 2(n-1)-1\}$
$u_{(2n-1)(n-3)+2(n-1)+1}$	v_{n-1}
$u_{(2n-1)(n-3)+2(n-1)+2}$	v_{n-1}
$u_{(2n-1)(n-3)+2(n-1)+3}$	v_0

Table 4.3: Definition of ${}_n\phi$





Figure 4.1: The bonding map ${}_3\phi$




Figure 4.2: The bonding map ${}_4\phi$




Figure 4.3: The bonding map 5ϕ

Let ${}_nK = \varprojlim \{{}_nX_i, {}_n\Phi_i^j\}_{i=0, j \geq i}^\infty$ be generated by $\{{}_n\phi, {}_n\phi, \dots\}$ (Definition 4.10).

Proposition 4.3. *Every proper nondegenerate subcontinuum of ${}_nK$ is an arc.*

Proof. Let W be a nondegenerate subcontinuum of ${}_nK$, and suppose W is not an arc. Since ${}_n\phi$ is simplicial, there exists $v \in V({}_nX_1)$ so that $v \in \pi_j(W)$ for infinitely many j . Then, $v_s \in \pi_j(W)$ for all j , where $s = 0$, $s = n$, or $s = n + 2$.

Case 1: Suppose $s = n$ or $s = n + 2$. Since W is not an arc and ${}_n\phi$ embeds $\langle u_{(2n-1)(n-3)+2(n-1)}, v_n, v_{n+2} \rangle$ onto $\langle v_0, v_n, v_{n+2} \rangle$, then $({}_nX_j \setminus ((\langle u_{(2n-1)(n-3)+2(n-1)}, v_n \rangle, {}_nX_j) \cup (\langle v_n, v_{n+2} \rangle, {}_nX_j))) \cap \pi_j(W) \neq \emptyset$ for infinitely many j (Definition 4.5), implying $(\langle u_{(2n-1)(n-3)+2(n-1)}, v_n \rangle, {}_nX_j) \subseteq \pi_j(W)$ for infinitely many j . Since ${}_n\Phi_0^2((\langle u_{(2n-1)(n-3)+2(n-1)}, v_n \rangle, {}_nX_2)) = {}_nX_0$, then ${}_nX_j \subseteq \pi_j(W)$ for all j , implying $W = {}_nK$.

Case 2: Suppose $s = 0$. Since W is not an arc and ${}_n\phi$ embeds $\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle$ onto $\langle v_0, v_{n-1} \rangle$, then $({}_nX_j \setminus ((\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle, {}_nX_j) \cap \pi_j(W) \neq \emptyset$ for infinitely many j . If $\pi_j(W) \cap (\bigcup_{i=1}^{n-3} (\langle v_0, u_i \rangle, {}_nX_j) \cup (\langle v_0, u_{(2n-1)(n-3)+1} \rangle, {}_nX_j) \cup (\langle v_0, u_{(2n-1)(n-3)+2(n-1)+1} \rangle, {}_nX_j) \setminus \{v_0\})) \neq \emptyset$ for some j , then $u_{(2n-1)(n-3)+2(n-1)+2} \in \pi_j(W)$. Thus, $(\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle, {}_nX_j) \subseteq \pi_j(W)$ for infinitely many j . Since ${}_n\Phi_0^4((\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle, {}_nX_4)) = {}_nX_0$, then ${}_nX_j \subseteq \pi_j(W)$ for each j , implying $W = {}_nK$. $\square$

Proposition 4.4. *${}_nK$ is atriodic.*

Proof. The claim follows from Proposition 4.3 and Theorem 3 in [6]. $\square$

Proposition 4.5. *Every proper nondegenerate subcontinuum of ${}_nK$ containing $(v_0, v_0, v_0, \dots)$ is an arc having $(v_0, v_0, v_0, \dots)$ as an endpoint.*

Proof. Let $W \subseteq {}_nK$ be a proper nondegenerate subcontinuum so that $(v_0, v_0, v_0, \dots) \in W$. By Proposition 4.3, W is an arc. Since W is proper and ${}_n\Phi_0^3((\langle v_0, v_{n-1} \rangle, {}_nX_3)) = {}_nX_0$, there exists a nonnegative integer N so that $v_{n-1} \notin \pi_j(W)$ for all $j \geq N$. Then, $\pi_j(W) \cap (\bigcup_{i=1}^{n-3} (\langle v_0, u_i \rangle, {}_nX_j) \cup (\langle v_0, u_{(2n-1)(n-3)+1} \rangle, {}_nX_j) \cup (\langle v_0, u_{(2n-1)(n-3)+2(n-1)+1} \rangle, {}_nX_j) \setminus \{v_0\}) = \emptyset$ for all $j \geq N$, implying $\pi_j(W) \subseteq (\langle v_0, v_{n-1} \rangle, {}_nX_j)$ for all $j \geq N$.

Since ${}_n\phi$ embeds $\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle$ onto $\langle v_0, v_{n-1} \rangle$ and ${}_n\phi(v_0) = v_0$, then $(v_0, v_0, \dots)$ is an endpoint of the arc W . $\square$

Let ${}_nS : V({}_nX_0) \longrightarrow \{\text{nonempty subsets of } E({}_nX_0)\}$ be an edge selection (Definition 4.7) on ${}_nX_0$ defined as: ${}_nS(v_i) = \{\langle v_0, v_i \rangle\}$ for all $i \in \{1, \dots, n\}$, ${}_nS(v_{n+1}) = \{\langle v_{n-1}, v_{n+1} \rangle\}$, ${}_nS(v_{n+2}) = \{\langle v_n, v_{n+2} \rangle\}$, and ${}_nS(v_0) = \{\langle v_0, v_i \rangle : i \in \{1, \dots, n-1\}\}$.

Proposition 4.6. ${}_n\phi$ preserves $({}_nS, {}_nS)$ (Definition 4.9).

Proof. i) (Recall Definition 4.5)

$$\begin{aligned}
 {}_n\phi((v_i, \langle v_0, v_i \rangle, {}_nX_1)) &= {}_n\phi(\langle u_{2(n-1)(n-3)+i}, v_i \rangle) = \langle v_n, v_{n+2} \rangle \in {}_nS(v_{n+2}) \\
 &= {}_nS({}_n\phi(v_i)) \text{ for all } i \in \{1, \dots, n-3\}, \\
 {}_n\phi((v_{n-2}, \langle v_0, v_{n-2} \rangle, {}_nX_1)) &= {}_n\phi(\langle u_{(2n-1)(n-3)+2(n-1)+1}, v_{n-2} \rangle) = \langle v_{n-1}, v_{n+1} \rangle \in \\
 &{}_nS(v_{n+1}) = {}_nS({}_n\phi(v_{n-2})), \\
 {}_n\phi((v_{n-1}, \langle v_0, v_{n-1} \rangle, {}_nX_1)) &= {}_n\phi(\langle u_{(2n-1)(n-3)+2(n-1)+3}, v_{n-1} \rangle) = \langle v_0, v_n \rangle \in {}_nS(v_n) \\
 &= {}_nS({}_n\phi(v_{n-1})), \\
 {}_n\phi((v_{n+1}, \langle v_{n-1}, v_{n+1} \rangle, {}_nX_1)) &= {}_n\phi(\langle v_{n-1}, v_{n+1} \rangle) = \langle v_n, v_{n+2} \rangle \in {}_nS(v_{n+2}) \\
 &= {}_nS({}_n\phi(v_{n+1})), \\
 {}_n\phi((v_n, \langle v_0, v_n \rangle, {}_nX_1)) &= {}_n\phi(\langle u_{(2n-1)(n-3)+2(n-1)}, v_n \rangle) = \langle v_0, v_n \rangle \in {}_nS(v_n) \\
 &= {}_nS({}_n\phi(v_n)), \\
 {}_n\phi((v_{n+2}, \langle v_n, v_{n+2} \rangle, {}_nX_1)) &= {}_n\phi(\langle v_n, v_{n+2} \rangle) = \langle v_n, v_{n+2} \rangle \in {}_nS(v_{n+2}) \\
 &= {}_nS({}_n\phi(v_{n+2})), \\
 {}_n\phi((v_0, \langle v_0, v_i \rangle, {}_nX_1)) &= {}_n\phi(\langle v_0, u_i \rangle) = \langle v_0, v_{n-1} \rangle \in {}_nS(v_0) \\
 &= {}_nS({}_n\phi(v_0)) \text{ for all } i \in \{1, \dots, n-3\}, \\
 {}_n\phi((v_0, \langle v_0, v_{n-2} \rangle, {}_nX_1)) &= {}_n\phi(\langle v_0, u_{(2n-1)(n-3)+2(n-1)+1} \rangle) = \langle v_0, v_{n-1} \rangle \in {}_nS(v_0) \\
 &= {}_nS({}_n\phi(v_0)), \text{ and} \\
 {}_n\phi((v_0, \langle v_0, v_{n-1} \rangle, {}_nX_1)) &= {}_n\phi(\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle) = \langle v_0, v_{n-1} \rangle \in {}_nS(v_0) \\
 &= {}_nS({}_n\phi(v_0)).
 \end{aligned}$$

ii)

$$\begin{aligned}
 {}_n\phi(\langle v_0, u_i \rangle) &= \langle v_0, v_{n-1} \rangle \in {}_nS(v_{n-1}) = {}_nS({}_n\phi(u_i)) \text{ for all } i \in \{1, \dots, n-3\}, \\
 {}_n\phi(\langle u_{(j-1)(n-3)+i}, u_{j(n-3)+i} \rangle) &= {}_n\phi(\langle u_{j(n-3)+i}, u_{(j+1)(n-3)+i} \rangle) = \langle v_0, v_{i-1+\frac{i}{2} \bmod n-2} \rangle \in \\
 &{}_nS(v_0) \cap {}_nS(v_{i-1+\frac{i}{2} \bmod n-2}) = {}_nS({}_n\phi(u_{(j-1)(n-3)+i})) \cap {}_nS({}_n\phi(u_{(j+1)(n-3)+i})) \cap \\
 &{}_nS({}_n\phi(u_{j(n-3)+i})) \text{ for all } i \in \{1, \dots, n-3\} \text{ and for all even} \\
 &j \in \{2, \dots, 2(n-1)-2\}, \\
 {}_n\phi(\langle u_{(2(n-1)-1)(n-3)+i}, u_{2(n-1)(n-3)+i} \rangle) &= \langle v_0, v_n \rangle \in {}_nS(v_n) = {}_nS({}_n\phi(u_{2(n-1)(n-3)+i})) \text{ for} \\
 &\text{all } i \in \{1, \dots, n-3\}, \\
 {}_n\phi(\langle v_0, u_{(2n-1)(n-3)+1} \rangle) &= \langle v_0, v_{n-1} \rangle \in {}_nS(v_{n-1}) = {}_nS({}_n\phi(u_{(2n-1)(n-3)+1})), \\
 {}_n\phi(\langle u_{(2n-1)(n-3)+2}, u_{(2n-1)(n-3)+3} \rangle) &= \langle v_0, v_{n-2} \rangle \in {}_nS(v_0) \cap {}_nS(v_{n-2}) = \\
 &{}_nS({}_n\phi(u_{(2n-1)(n-3)+2})) \cap {}_nS({}_n\phi(u_{(2n-1)(n-3)+3})), \\
 {}_n\phi(\langle u_{(2n-1)(n-3)+i-1}, u_{(2n-1)(n-3)+i} \rangle) &= {}_n\phi(\langle u_{(2n-1)(n-3)+i}, u_{(2n-1)(n-3)+i+1} \rangle) = \langle v_0, v_{\frac{i-3}{2}} \rangle \\
 &\in {}_nS(v_0) \cap {}_nS(v_{\frac{i-3}{2}}) = {}_nS({}_n\phi(u_{(2n-1)(n-3)+i-1})) \cap {}_nS({}_n\phi(u_{(2n-1)(n-3)+i+1})) \cap \\
 &{}_nS({}_n\phi(u_{(2n-1)(n-3)+i})) \text{ for all odd } i \in \{5, \dots, 2(n-1)-1\}, \\
 {}_n\phi(\langle v_0, u_{(2n-1)(n-3)+2(n-1)+1} \rangle) &= \langle v_0, v_{n-1} \rangle \in {}_nS(v_{n-1}) = {}_nS({}_n\phi(u_{(2n-1)(n-3)+2(n-1)+1})), \\
 &\text{and} \\
 {}_n\phi(\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle) &= {}_n\phi(\langle u_{(2n-1)(n-3)+2(n-1)+2}, u_{(2n-1)(n-3)+2(n-1)+3} \rangle) = \\
 &\langle v_0, v_{n-1} \rangle \in {}_nS(v_0) \cap {}_nS(v_{n-1}) = {}_nS({}_n\phi(u_{(2n-1)(n-3)+2(n-1)+3})) \cap \\
 &{}_nS({}_n\phi(u_{(2n-1)(n-3)+2(n-1)+2})).
 \end{aligned}$$

With (i) above, condition (ii) of Definition 4.9 is satisfied whenever $v \in V({}_nX_0)$. $\square$

Let $D({}_nX_0)$ be as in Definition 4.1. Then, $V(D({}_nX_0)) = \{a_i : i \in \{1, \dots, n+2\}\}$ and $E(D({}_nX_0)) = \{\langle a_i, a_j \rangle : i, j \in \{1, \dots, n\} \text{ and } i \neq j\} \cup \{\langle a_{n-1}, a_{n+1} \rangle, \langle a_n, a_{n+2} \rangle\}$, where $a_i^* = \langle v_0, v_i \rangle$ for all $i \in \{1, \dots, n\}$ and $a_{n+1}^* = \langle v_{n-1}, v_{n+1} \rangle$ and $a_{n+2}^* = \langle v_n, v_{n+2} \rangle$.

Let $D({}_n\phi, {}_nX_1)$ be as in Definition 4.2. Then, $V(D({}_n\phi, {}_nX_1)) = \{b_i : i \in \{0, \dots, n(n-3)+n+3\}\}$ and $E(D({}_n\phi, {}_nX_1)) = \{\langle b_0, b_i \rangle : i \in \{1, \dots, n-3\} \cup \{n(n-3)+1, n(n-3)+n+1, n(n-3)+n+2\}\} \cup \{\langle b_{j(n-3)+i}, b_{(j+1)(n-3)+i} \rangle : i \in \{1, \dots, n-3\} \text{ and } j \in \{0, \dots, n-2\}\} \cup \{\langle b_{n(n-3)+i}, b_{n(n-3)+i+1} \rangle : i \in \{1, \dots, n-1\}\} \cup \{\langle b_{n(n-3)+n+2},$

$b_{n(n-3)+n+3}\rangle\}$, where b^* for $b \in V(D(n\phi, nX_1))$ is given in the following table.

$$\begin{aligned}
 b_0^* &= \bigcup_{i=1}^{n-3} \langle v_0, u_i \rangle \bigcup_{i=1}^{n-3} \langle u_i, u_{n-3+i} \rangle \bigcup \langle v_0, u_{(2n-1)(n-3)+1} \rangle \bigcup \langle u_{(2n-1)(n-3)+1}, \\
 &\quad u_{(2n-1)(n-3)+2} \rangle \bigcup \langle v_0, u_{(2n-1)(n-3)+2(n-1)+1} \rangle \bigcup \langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle \\
 &\quad \bigcup \langle u_{(2n-1)(n-3)+2(n-1)+2}, u_{(2n-1)(n-3)+2(n-1)+3} \rangle \\
 b_{j(n-3)+i}^* &= \langle u_{(2j+1)(n-3)+i}, u_{(2j+2)(n-3)+i} \rangle \bigcup \langle u_{(2j+2)(n-3)+i}, u_{(2j+3)(n-3)+i} \rangle \\
 &\quad \text{for each } i \in \{1, \dots, n-3\} \text{ and for each } j \in \{0, \dots, n-3\} \\
 b_{(n-2)(n-3)+i}^* &= \langle u_{(2(n-1)-1)(n-3)+i}, u_{2(n-1)(n-3)+i} \rangle \text{ for each } i \in \{1, \dots, n-3\} \\
 b_{(n-1)(n-3)+i}^* &= \langle u_{2(n-1)(n-3)+i}, v_i \rangle \text{ for each } i \in \{1, \dots, n-3\} \\
 b_{n(n-3)+i}^* &= \langle u_{(2n-1)(n-3)+2i}, u_{(2n-1)(n-3)+2i+1} \rangle \bigcup \langle u_{(2n-1)(n-3)+2i+1}, u_{(2n-1)(n-3)+2i+2} \rangle \\
 &\quad \text{for each } i \in \{1, \dots, n-2\} \\
 b_{n(n-3)+n-1}^* &= \langle u_{(2n-1)(n-3)+2(n-1)}, v_n \rangle \\
 b_{n(n-3)+n}^* &= \langle v_n, v_{n+2} \rangle \\
 b_{n(n-3)+n+1}^* &= \langle u_{(2n-1)(n-3)+2(n-1)+1}, v_{n-2} \rangle \\
 b_{n(n-3)+n+2}^* &= \langle u_{(2n-1)(n-3)+2(n-1)+3}, v_{n-1} \rangle \\
 b_{n(n-3)+n+3}^* &= \langle v_{n-1}, v_{n+1} \rangle
 \end{aligned}$$

Table 4.4: Subgraphs of nX_1 corresponding to vertices of $D(n\phi, nX_1)$

Let $B^i \subseteq D(n\phi, nX_1)$ for each $i \in \{1, \dots, n\}$ be the arc given by the following table.

arc	as a subgraph
B^i	$\langle b_0, b_i, b_{n-3+i}, b_{2(n-3)+i}, \dots, b_{(n-1)(n-3)+i} \rangle$ for all $i \in \{1, \dots, n-3\}$
B^{n-2}	$\langle b_0, b_{n(n-3)+n+1} \rangle$
B^{n-1}	$\langle b_0, b_{n(n-3)+n+2}, b_{n(n-3)+n+3} \rangle$
B^n	$\langle b_0, b_{n(n-3)+1}, \dots, b_{n(n-3)+n} \rangle$

Table 4.5: Determination of $D(n\phi, nX_1)$

Then, $\bigcup_{i=1}^n B^i = D(n\phi, nX_1)$ and $B^i \cap B^j = \{b_0\}$ for all distinct $i, j \in \{1, \dots, n\}$.

Proposition 4.7. *Let $d[n\phi]$ be as in Definition 4.2. Then, $d[n\phi]$ is an ultra light simplicial map (Definition 4.4).*

Proof. $d[n\phi] : D(n\phi, nX_1) \longrightarrow D(nX_0)$, where $b \in V(D(n\phi, nX_1))$, and the subgraphs forming the inverse images of edges under $d[n\phi]$ are given in the next tables.

b	$d[n\phi](b)$
b_0	a_{n-1}
$b_{j(n-3)+i}$	$a_{i-1+\frac{2i+2}{2} \bmod n-2} = a_{i+j \bmod n-2}$ for all $i \in \{1, \dots, n-3\}$ and for all $j \in \{0, \dots, n-3\}$
$b_{(n-2)(n-3)+i}$	a_n for all $i \in \{1, \dots, n-3\}$
$b_{(n-1)(n-3)+i}$	a_{n+2} for all $i \in \{1, \dots, n-3\}$
$b_{n(n-3)+1}$	a_{n-2}
$b_{n(n-3)+i}$	$a_{\frac{2i+1-3}{2}} = a_{i-1}$ for all $i \in \{2, \dots, n-2\}$
$b_{n(n-3)+n-1}$	a_n
$b_{n(n-3)+n}$	a_{n+2}
$b_{n(n-3)+n+1}$	a_{n+1}
$b_{n(n-3)+n+2}$	a_n
$b_{n(n-3)+n+3}$	a_{n+2}

Table 4.6: Determination of $d[n\phi]$




Figure 4.4: The map $d[5\phi]$

$$\begin{aligned}
 d[n\phi]^{-1}(\langle a_{n-1}, a_i \rangle) &= \langle b_0, b_i \rangle \text{ for all } i \in \{1, \dots, n-3\} \\
 d[n\phi]^{-1}(\langle a_{n-1}, a_{n-2} \rangle) &= \langle b_0, b_{n(n-3)+1} \rangle \\
 d[n\phi]^{-1}(\langle a_{n-1}, a_{n+1} \rangle) &= \langle b_0, b_{n(n-3)+n+1} \rangle \\
 d[n\phi]^{-1}(\langle a_{n-1}, a_n \rangle) &= \langle b_0, b_{n(n-3)+n+2} \rangle \\
 d[n\phi]^{-1}(\langle a_i, a_{i+1} \rangle) &= \bigcup_{j=0}^{i-1} \langle b_{j(n-3)+i-j}, b_{(j+1)(n-3)+i-j} \rangle \bigcup_{j=1}^{n-4-i} \langle b_{(n-3-j)(n-3)+i+1+j}, \\
 &\quad b_{(n-2-j)(n-3)+i+1+j} \rangle \bigcup \langle b_{n(n-3)+i+1}, b_{n(n-3)+i+2} \rangle \\
 &\quad \text{for all } i \in \{1, \dots, n-4\} \\
 d[n\phi]^{-1}(\langle a_{n-3}, a_{n-2} \rangle) &= \bigcup_{i=1}^{n-3} \langle b_{(n-3-i)(n-3)+i}, b_{(n-2-i)(n-3)+i} \rangle \\
 d[n\phi]^{-1}(\langle a_i, a_n \rangle) &= \langle b_{(n-3)(n-3)+i+1}, b_{(n-2)(n-3)+i+1} \rangle \text{ for all } i \in \{1, \dots, n-4\} \\
 d[n\phi]^{-1}(\langle a_{n-3}, a_n \rangle) &= \langle b_{n(n-3)+n-2}, b_{n(n-3)+n-1} \rangle \\
 d[n\phi]^{-1}(\langle a_{n-2}, a_n \rangle) &= \langle b_{(n-3)(n-3)+1}, b_{(n-2)(n-3)+1} \rangle \\
 d[n\phi]^{-1}(\langle a_n, a_{n+2} \rangle) &= \langle b_{n(n-3)+n-1}, b_{n(n-3)+n} \rangle \bigcup \langle b_{n(n-3)+n+2}, b_{n(n-3)+n+3} \rangle \\
 &\quad \bigcup_{i=1}^{n-3} \langle b_{(n-2)(n-3)+i}, b_{(n-1)(n-3)+i} \rangle
 \end{aligned}$$

 Table 4.7: Edge inverses under $d[n\phi]$

Each subgraph is a union of disjoint edges, and so each component of an edge inverse mapping onto that edge is an edge. $\square$

Let ${}_nY_1$ be a subdivision of ${}_nX_0$, where ${}_nB_1^i$ is a subdivision of ${}_nA_0^i$ for each $i \in \{1, \dots, n\}$ defined in the following table.

arc	as a subgraph
${}_nB_1^i$	$\langle v_0, w_i, w_{n-3+i}, w_{2(n-3)+i}, \dots, w_{(n-2)(n-3)+i}, v_i \rangle$ for all $i \in \{1, \dots, n-3\}$
${}_nB_1^{n-2}$	$\langle v_0, v_{n-2} \rangle = {}_nA_0^{n-2}$
${}_nB_1^{n-1}$	$\langle v_0, v_{n-1}, v_{n+1} \rangle = {}_nA_0^{n-1}$
${}_nB_1^n$	$\langle v_0, w_{(n-1)(n-3)+1}, \dots, w_{(n-1)(n-3)+n-2}, v_n, v_{n+2} \rangle$

 Table 4.8: Definition of ${}_nY_1$

Define ${}_n\lambda : {}_nY_1 \longrightarrow D({}_n\phi, {}_nX_1)$ to be the simplicial isomorphism determined by the following table where $v \in V({}_nY_1)$.

v	${}_n\lambda(v)$
v_0	b_0
v_i	$b_{(n-1)(n-3)+i}$ for all $i \in \{1, \dots, n-3\}$
v_{n-2}	$b_{n(n-3)+n+1}$
v_{n-1}	$b_{n(n-3)+n+2}$
v_n	$b_{n(n-3)+n-1}$
v_{n+1}	$b_{n(n-3)+n+3}$
v_{n+2}	$b_{n(n-3)+n}$
$w_{j(n-3)+i}$	$b_{j(n-3)+i}$ for all $j \in \{0, \dots, n-2\}$ and for all $i \in \{1, \dots, n-3\}$
$w_{(n-1)(n-3)+i}$	$b_{n(n-3)+i}$ for all $i \in \{1, \dots, n-2\}$

Table 4.9: Definition of ${}_n\lambda$

Proposition 4.8. ${}_n\lambda$ is a consistency isomorphism (Definition 4.8).

Proof. i)

$$\begin{aligned}
 (v_i, \langle v_0, v_i \rangle, {}_nX_1) &= \langle u_{2(n-1)(n-3)+i}, v_i \rangle = b_{(n-1)(n-3)+i}^* \\
 &= [{}_n\lambda(v_i)]^* \text{ for all } i \in \{1, \dots, n-3\}, \\
 (v_{n-2}, \langle v_0, v_{n-2} \rangle, {}_nX_1) &= \langle u_{(2n-1)(n-3)+2(n-1)+1}, v_{n-2} \rangle = b_{n(n-3)+n+1}^* \\
 &= [{}_n\lambda(v_{n-2})]^*, \\
 (v_{n-1}, \langle v_0, v_{n-1} \rangle, {}_nX_1) &= \langle u_{(2n-1)(n-3)+2(n-1)+3}, v_{n-1} \rangle = b_{n(n-3)+n+2}^* \\
 &= [{}_n\lambda(v_{n-1})]^*, \\
 (v_{n+1}, \langle v_{n-1}, v_{n+1} \rangle, {}_nX_1) &= \langle v_{n-1}, v_{n+1} \rangle = b_{n(n-3)+n+3}^* \\
 &= [{}_n\lambda(v_{n+1})]^*, \\
 (v_n, \langle v_0, v_n \rangle, {}_nX_1) &= \langle u_{(2n-1)(n-3)+2(n-1)}, v_n \rangle = b_{n(n-3)+n-1}^* \\
 &= [{}_n\lambda(v_n)]^*,
 \end{aligned}$$

$$\begin{aligned}
 (v_{n+2}, \langle v_n, v_{n+2} \rangle, {}_nX_1) &= \langle v_n, v_{n+2} \rangle = b_{n(n-3)+n}^* \\
 &= [{}_n\lambda(v_{n+2})]^*, \\
 (v_0, \langle v_0, v_i \rangle, {}_nX_1) &= \langle v_0, v_i \rangle = b_0^* \\
 &= [{}_n\lambda(v_0)]^* \text{ for all } i \in \{1, \dots, n-3\}, \\
 (v_0, \langle v_0, v_{n-2} \rangle, {}_nX_1) &= \langle v_0, u_{(2n-1)(n-3)+2(n-1)+1} \rangle = b_0^* \\
 &= [{}_n\lambda(v_0)]^*, \text{ and} \\
 (v_0, \langle v_0, v_{n-1} \rangle, {}_nX_1) &= \langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle = b_0^* \\
 &= [{}_n\lambda(v_0)]^*.
 \end{aligned}$$

ii)

$$\begin{aligned}
 [{}_n\lambda(w_{j(n-3)+i})]^* &= b_{j(n-3)+i}^* = \langle u_{(2j+1)(n-3)+i}, u_{(2j+2)(n-3)+i} \rangle \cup \langle u_{(2j+2)(n-3)+i}, \\
 &\quad u_{(2j+3)(n-3)+i} \rangle \subseteq {}_nA_1^i = (\langle v_0, v_i \rangle, {}_nX_1) \text{ for all } i \in \{1, \dots, n-3\} \text{ and} \\
 &\quad \text{for all } j \in \{0, \dots, n-3\}, \\
 [{}_n\lambda(w_{(n-2)(n-3)+i})]^* &= b_{(n-2)(n-3)+i}^* = \langle u_{(2(n-1)-1)(n-3)+i}, u_{2(n-1)(n-3)+i} \rangle \subseteq {}_nA_1^i \\
 &= (\langle v_0, v_i \rangle, {}_nX_1) \text{ for all } i \in \{1, \dots, n-3\}, \text{ and} \\
 [{}_n\lambda(w_{(n-1)(n-3)+i})]^* &= b_{n(n-3)+i}^* = \langle u_{(2n-1)(n-3)+2i}, u_{(2n-1)(n-3)+2i+1} \rangle \cup \\
 &\quad \langle u_{(2n-1)(n-3)+2i+1}, u_{(2n-1)(n-3)+2i+2} \rangle \subseteq {}_nA_1^i = (\langle v_0, v_n \rangle, {}_nX_1) \\
 &\quad \text{for each } i \in \{1, \dots, n-2\}. \quad \square
 \end{aligned}$$

Proposition 4.9. ${}_n\phi$ is consistent on ${}_nS$ (Definition 4.7).

Proof. The claim follows from Proposition 4.8 and Definition 4.7. $\square$

Let $f : G_1 \longrightarrow G_0$ be a light simplicial map between graphs and $x_0, x_1, \dots, x_m$ for some $m \in \{1, 2, \dots\}$ be vertices of G_1 so that x_i and x_{i+1} are adjacent for each $i \in \{0, \dots, m-1\}$. Define $\bar{f}(\langle x_0, x_1, \dots, x_m \rangle) = \langle f(x_0), f(x_1), \dots, f(x_m) \rangle$.

For each $j \in \{0, 1, \dots\}$, let ${}_nA_j^i = (\langle v_0, v_i \rangle, {}_nX_j)$ for each $i \in \{1, \dots, n-2\}$, ${}_nA_j^{n-1} = (\langle v_0, v_{n-1} \rangle, {}_nX_j) \vee (\langle v_{n-1}, v_{n+1} \rangle, {}_nX_j)$, and ${}_nA_j^n = (\langle v_0, v_n \rangle, {}_nX_j) \vee (\langle v_n, v_{n+2} \rangle, {}_nX_j)$ (Definition 4.5).

Proposition 4.10. *Let $m \in \{1, 2, \dots\}$ and, for each $i \in \{1, \dots, n\}$, $\langle x_0^i, x_1^i, \dots, x_{k_i}^i \rangle = \overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^i)$ for some $k_i \in \{1, 2, \dots\}$. Then,*

$$i) \overline{{}_n\Phi_0^m}({}_nA_m^i) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle \vee \langle x_{k_{n-1}-1}^{n-1}, x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle \bigvee_{j=0}^{n-3} (\langle x_0^{i+j \bmod n-2}, x_1^{i+j \bmod n-2}, \dots, x_{k_{i+j \bmod n-2}}^{i+j \bmod n-2} \rangle \vee \langle x_{k_{i+j \bmod n-2}-1}^{i+j \bmod n-2}, \dots, x_0^{i+j \bmod n-2} \rangle) \vee \overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^n) \text{ for each } i \in \{1, \dots, n-3\},$$

$$ii) \overline{{}_n\Phi_0^m}({}_nA_m^{n-2}) = \overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^{n-1}),$$

$$iii) \overline{{}_n\Phi_0^m}({}_nA_m^{n-1}) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle \vee \langle x_{k_{n-1}-1}^{n-1}, x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle \vee \overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^n), \text{ and}$$

$$iv) \overline{{}_n\Phi_0^m}({}_nA_m^n) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle \vee \langle x_{k_{n-1}-1}^{n-1}, x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle \vee \langle x_0^{n-2}, x_1^{n-2}, \dots, x_{k_{n-2}}^{n-2} \rangle \vee \langle x_{k_{n-2}-1}^{n-2}, \dots, x_0^{n-2} \rangle \bigvee_{j=1}^{n-3} (\langle x_0^j, x_1^j, \dots, x_{k_j}^j \rangle \vee \langle x_{k_j}^j, x_{k_j-1}^j, \dots, x_0^j \rangle) \vee \overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^n).$$

Proof. **CLAIM:** $(\langle v_{n-1}, v_n \rangle, {}_nX_i) = \langle v_{n-1}, v_{n+1} \rangle$ for each $i \in \{1, 2, \dots\}$. Since ${}_n\Phi_{i-1}^i$ is a subdivision of ${}_n\phi$ matching ${}_nX_{i-1}$ for each $i \in \{1, 2, \dots\}$ and ${}_n\phi$ embeds $\langle v_n, v_{n+2} \rangle$ onto $\langle v_n, v_{n+2} \rangle$, then $(\langle v_n, v_{n+2} \rangle, {}_nX_i) = \langle v_n, v_{n+2} \rangle$ for each $i \in \{1, 2, \dots\}$. Since ${}_n\phi$ embeds $\langle v_{n-1}, v_{n+1} \rangle$ onto $\langle v_n, v_{n+2} \rangle$, then $(\langle v_{n-1}, v_{n+1} \rangle, {}_nX_i) = \langle v_{n-1}, v_{n+1} \rangle$ for each $i \in \{1, 2, \dots\}$.

i) Since ${}_n\Phi_{m-1}^m$ is a subdivision of ${}_n\phi$ matching ${}_nX_{m-1}$ and ${}_n\phi$ embeds $\langle v_0, u_i \rangle$ onto $\langle v_0, v_{n-1} \rangle, \langle u_i, u_{n-3+i} \rangle$ onto $\langle v_{n-1}, v_0 \rangle, \langle u_{j(n-3)+i}, u_{(j+1)(n-3)+i} \rangle$ onto $\langle v_0, v_{i-1+\frac{i+1}{2} \bmod n-2} \rangle$ and $\langle u_{(j+1)(n-3)+i}, u_{(j+2)(n-3)+i} \rangle$ onto $\langle v_{i-1+\frac{i+1}{2} \bmod n-2}, v_0 \rangle$ for each odd $j \in \{1, \dots, 2(n-1)-3\}$, and $\langle u_{(2(n-1)-1)(n-3)+i}, u_{2(n-1)(n-3)+i}, v_i \rangle$ onto $\langle v_0, v_n, v_{n+2} \rangle$, for each $i \in \{1, \dots, n-3\}$, then $\overline{{}_n\Phi_{m-1}^m}({}_nA_{m-1}^i) = (\langle v_0, v_{n-1} \rangle, {}_nX_{m-1}) \vee (\langle v_{n-1}, v_0 \rangle, {}_nX_{m-1}) \bigvee_{j=0}^{n-3} ((\langle v_0, v_{i+j \bmod n-2} \rangle, {}_nX_{m-1}) \vee (\langle v_{i+j \bmod n-2}, v_0 \rangle, {}_nX_{m-1})) \vee {}_nA_{m-1}^n$ for each $i \in \{1, \dots, n-3\}$. By the claim, $\overline{{}_n\Phi_0^{m-1}}((\langle v_0, v_{n-1} \rangle, {}_nX_{m-1})) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle$. Thus, $\overline{{}_n\Phi_0^m}({}_nA_m^i) = \overline{{}_n\Phi_0^{m-1}}(\overline{{}_n\Phi_{m-1}^m}({}_nA_{m-1}^i)) = \overline{{}_n\Phi_0^{m-1}}((\langle v_0, v_{n-1} \rangle, {}_nX_{m-1}) \vee (\langle v_{n-1}, v_0 \rangle, {}_nX_{m-1}) \bigvee_{j=0}^{n-3} ((\langle v_0, v_{i+j \bmod n-2} \rangle, {}_nX_{m-1}) \vee (\langle v_{i+j \bmod n-2}, v_0 \rangle, {}_nX_{m-1})) \vee {}_nA_{m-1}^n) = \overline{{}_n\Phi_0^{m-1}}((\langle v_0, v_{n-1} \rangle, {}_nX_{m-1}) \vee (\langle v_{n-1}, v_0 \rangle, {}_nX_{m-1}) \bigvee_{j=0}^{n-3} (\overline{{}_n\Phi_0^{m-1}}((\langle v_{n-1}, v_0 \rangle, {}_nX_{m-1})) \vee \overline{{}_n\Phi_0^{m-1}}((\langle v_0, v_{i+j \bmod n-2} \rangle, {}_nX_{m-1})) \vee \overline{{}_n\Phi_0^{m-1}}((\langle v_{i+j \bmod n-2}, v_0 \rangle, {}_nX_{m-1}))) \vee \overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^n)) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle \vee \langle x_{k_{n-1}-1}^{n-1}, x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle$

$x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle \bigvee_{j=0}^{n-3} (\langle x_0^{i+j \bmod n-2}, x_1^{i+j \bmod n-2}, \dots, x_{k_{i+j \bmod n-2}}^{i+j \bmod n-2} \rangle \vee \langle x_{k_{i+j \bmod n-2}}^{i+j \bmod n-2}, x_{k_{i+j \bmod n-2}-1}^{i+j \bmod n-2}, \dots, x_0^{i+j \bmod n-2} \rangle) \vee \overline{n\Phi_0^{m-1}}(nA_{m-1}^n)$ for each $i \in \{1, \dots, n-3\}$.

ii) Since $n\Phi_{m-1}^m$ is a subdivision of $n\phi$ matching nX_{m-1} and $n\phi$ embeds $\langle v_0, u_{(2n-1)(n-3)+2(n-1)+1}, v_{n-2} \rangle$ onto $\langle v_0, v_{n-1}, v_{n+1} \rangle$, then $\overline{n\Phi_{m-1}^m}(nA_m^{n-2}) = nA_{m-1}^{n-1}$, giving $\overline{n\Phi_0^m}(nA_m^{n-2}) = \overline{n\Phi_0^{m-1}}(\overline{n\Phi_{m-1}^m}(nA_m^{n-2})) = \overline{n\Phi_0^{m-1}}(nA_{m-1}^{n-1})$.

iii) Since $n\Phi_{m-1}^m$ is a subdivision of $n\phi$ matching nX_{m-1} and $n\phi$ embeds $\langle v_0, u_{(2n-1)(n-3)+2(n-1)+2} \rangle$ onto $\langle v_0, v_{n-1} \rangle$, $\langle u_{(2n-1)(n-3)+2(n-1)+2}, u_{(2n-1)(n-3)+2(n-1)+3} \rangle$ onto $\langle v_{n-1}, v_0 \rangle$, and $\langle u_{(2n-1)(n-3)+2(n-1)+3}, v_{n-1}, v_{n+1} \rangle$ onto $\langle v_0, v_n, v_{n+2} \rangle$, then $\overline{n\Phi_{m-1}^m}(nA_m^{n-1}) = (\langle v_0, v_{n-1} \rangle, nX_{m-1}) \vee (\langle v_{n-1}, v_0 \rangle, nX_{m-1}) \vee nA_{m-1}^n$, and so by the claim, $\overline{n\Phi_0^m}(nA_m^{n-1}) = \overline{n\Phi_0^{m-1}}(\overline{n\Phi_{m-1}^m}(nA_m^{n-1})) = \overline{n\Phi_0^{m-1}}((\langle v_0, v_{n-1} \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}((\langle v_{n-1}, v_0 \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}(nA_{m-1}^n) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle \vee \langle x_{k_{n-1}-1}^{n-1}, x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle \vee \overline{n\Phi_0^{m-1}}(nA_{m-1}^n)$.

iv) Since $n\Phi_{m-1}^m$ is a subdivision of $n\phi$ matching nX_{m-1} and $n\phi$ embeds $\langle v_0, u_{(2n-1)(n-3)+1} \rangle$ onto $\langle v_0, v_{n-1} \rangle$, $\langle u_{(2n-1)(n-3)+1}, u_{(2n-1)(n-3)+2} \rangle$ onto $\langle v_{n-1}, v_0 \rangle$, $\langle u_{(2n-1)(n-3)+2}, u_{(2n-1)(n-3)+3} \rangle$ onto $\langle v_0, v_{n-2} \rangle$, $\langle u_{(2n-1)(n-3)+3}, u_{(2n-1)(n-3)+4} \rangle$ onto $\langle v_{n-2}, v_0 \rangle$, $\langle u_{(2n-1)(n-3)+i}, u_{(2n-1)(n-3)+i+1} \rangle$ onto $\langle v_0, v_{\frac{i-2}{2}} \rangle$ and $\langle u_{(2n-1)(n-3)+i+1}, u_{(2n-1)(n-3)+i+2} \rangle$ onto $\langle v_{\frac{i-2}{2}}, v_0 \rangle$ for each even $i \in \{4, \dots, 2(n-1)-2\}$, and $\langle u_{((2n-1)(n-3)+2(n-1))(n-3)+i}, v_n, v_{n+2} \rangle$ onto $\langle v_0, v_n, v_{n+2} \rangle$, then $\overline{n\Phi_{m-1}^m}(nA_m^n) = (\langle v_0, v_{n-1} \rangle, nX_{m-1}) \vee (\langle v_{n-1}, v_0 \rangle, nX_{m-1}) \vee (\langle v_0, v_{n-2} \rangle, nX_{m-1}) \vee (\langle v_{n-2}, v_0 \rangle, nX_{m-1}) \bigvee_{i=1}^{n-3} ((\langle v_0, v_i \rangle, nX_{m-1}) \vee (\langle v_i, v_0 \rangle, nX_{m-1})) \vee nA_{m-1}^n$, and so by the claim, $\overline{n\Phi_0^m}(nA_m^n) = \overline{n\Phi_0^{m-1}}(\overline{n\Phi_{m-1}^m}(nA_m^n)) = \overline{n\Phi_0^{m-1}}((\langle v_0, v_{n-1} \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}((\langle v_{n-1}, v_0 \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}((\langle v_0, v_{n-2} \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}((\langle v_{n-2}, v_0 \rangle, nX_{m-1})) \bigvee_{i=1}^{n-3} \overline{n\Phi_0^{m-1}}((\langle v_0, v_i \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}((\langle v_i, v_0 \rangle, nX_{m-1})) \vee \overline{n\Phi_0^{m-1}}(nA_{m-1}^n) = \langle x_0^{n-1}, x_1^{n-1}, \dots, x_{k_{n-1}-1}^{n-1} \rangle \vee \langle x_{k_{n-1}-1}^{n-1}, x_{k_{n-1}-2}^{n-1}, \dots, x_0^{n-1} \rangle \vee \langle x_0^{n-2}, x_1^{n-2}, \dots, x_{k_{n-2}}^{n-2} \rangle \vee \langle x_{k_{n-2}}^{n-2}, x_{k_{n-2}-1}^{n-2}, \dots, x_0^{n-2} \rangle \bigvee_{i=1}^{n-3} (\langle x_0^i, x_1^i, \dots, x_{k_i}^i \rangle \vee \langle x_{k_i}^i, x_{k_i-1}^i, \dots, x_0^i \rangle) \vee \overline{n\Phi_0^{m-1}}(nA_{m-1}^n)$.

□

Let $p : \{1, \dots, n-2\} \longrightarrow \{1, \dots, n-2\}$ be the bijection defined as $p(i) = i+1$

for each $i \in \{1, \dots, n-3\}$ and $p(n-2) = 1$.

Let $C = \{\langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle, \langle v_0, v_{n-1}, v_0, v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{p^j(n-2)}, v_0, \underline{v_{p^j(n-3)}, v_0}, \dots, \underline{v_{p^j(1)}, v_0}, v_{n-1}, v_0 \rangle, \langle v_0, v_{n-1}, v_{n+1}, v_{n-1}, v_0 \rangle, \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle, \langle v_0, v_{n-1}, v_0, v_n, v_{n+2}, v_n, v_0, v_{n-1}, v_0 \rangle : j \in \{1, \dots, n-2\}\}$.

Proposition 4.11. *Let $e \in C$, $k \in \{1, 2, \dots\}$, and $\{x_0, x_1, \dots, x_k\} \subseteq V(nX_0)$ so that $e = \langle x_0, x_1, \dots, x_k \rangle$. Then, $\langle x_k, x_{k-1}, \dots, x_0 \rangle = e$.*

Proof. If $v_{n+1} \in e$, then v_{n+1} is a vertex of symmetry for e . If $v_{n+2} \in e$, then v_{n+2} is a vertex of symmetry for e . Otherwise, v_n is a vertex of symmetry for e . $\square$

Proposition 4.12. *Let $m \geq 3$. Then, $\overline{{}_n\Phi_0^{m-1}}({}_nA_{m-1}^i) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{i,m-1} \vee e_2^{i,m-1} \vee \dots \vee e_{k_{i,m-1}}^{i,m-1} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle$ for some $k_{i,m-1} \in \{0, 1, \dots\}$ for each $i \in \{1, \dots, n\} \setminus \{n-2\}$ and $\overline{{}_n\Phi_0^m}({}_nA_m^{n-2}) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{n-2,m} \vee e_2^{n-2,m} \vee \dots \vee e_{k_{n-2,m}}^{n-2,m} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle$ for some $k_{n-2,m} \in \{0, 1, \dots\}$, where $e_k^{i,m-1} \in C$ for each $k \in \{1, \dots, k_{i,m-1}\}$ and each $i \in \{1, \dots, n\} \setminus \{n-2\}$ and $e_k^{n-2,m} \in C$ for each $k \in \{1, \dots, k_{n-2,m}\}$.*

Proof. By Proposition 4.10, $\overline{{}_n\Phi_0^0}({}_nA_0^i) = \langle v_0, v_i \rangle$ for all $i \in \{1, \dots, n-2\}$, $\overline{{}_n\Phi_0^0}({}_nA_0^{n-1}) = \langle v_0, v_{n-1}, v_{n+1} \rangle$, $\overline{{}_n\Phi_0^0}({}_nA_0^n) = \langle v_0, v_n, v_{n+2} \rangle$; $\overline{{}_n\Phi_0^1}({}_nA_1^i) = \langle v_0, v_{n-1}, v_0, v_{p^{i-1}(1)}, v_0, \underline{v_{p^{i-1}(2)}, v_0}, \dots, \underline{v_{p^{i-1}(n-2)}, v_0}, v_n, v_{n+2} \rangle$ for all $i \in \{1, \dots, n-3\}$, $\overline{{}_n\Phi_0^1}({}_nA_1^{n-2}) = \langle v_0, v_{n-1}, v_{n+1} \rangle$, $\overline{{}_n\Phi_0^1}({}_nA_1^{n-1}) = \langle v_0, v_{n-1}, v_0, v_n, v_{n+2} \rangle$, $\overline{{}_n\Phi_0^1}({}_nA_1^n) = \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle$; $\overline{{}_n\Phi_0^2}({}_nA_2^i) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{j=i-1}^{n-4} \langle v_0, v_{n-1}, v_0, v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{p^j(n-2)}, v_0, \underline{v_{p^j(n-3)}, v_0}, \dots, \underline{v_{p^j(1)}, v_0}, v_{n-1}, v_0 \rangle \vee \langle v_0, v_{n-1}, v_{n+1}, v_{n-1}, v_0 \rangle \bigvee_{j=0}^{i-2} \langle v_0, v_{n-1}, v_0, v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{p^j(n-2)}, v_0, \underline{v_{p^j(n-3)}, v_0}, \dots, \underline{v_{p^j(1)}, v_0}, v_{n-1}, v_0 \rangle \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle$ for all $i \in \{1, \dots, n-3\}$, $\overline{{}_n\Phi_0^2}({}_nA_2^{n-2}) = \langle v_0, v_{n-1}, v_0, v_n, v_{n+2} \rangle$, $\overline{{}_n\Phi_0^2}({}_nA_2^{n-1}) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle$, $\overline{{}_n\Phi_0^2}({}_nA_2^n) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee \langle v_0,$

$$\begin{aligned} & \langle v_{n-1}, v_{n+1}, v_{n-1}, v_0 \rangle \bigvee_{i=1}^{n-3} \langle v_0, v_{n-1}, v_0, v_{p^{i-1}(1)}, v_0, \underline{v_{p^{i-1}(2)}, v_0}, \dots, \underline{v_{p^{i-1}(n-2)}, v_0}}, v_n, v_{n+2}, v_n, \\ & v_0, v_{p^{i-1}(n-2)}, v_0, \underline{v_{p^{i-1}(n-3)}, v_0}, \dots, \underline{v_{p^{i-1}(1)}, v_0}}, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \\ & \underline{v_0}, \dots, \underline{v_{n-3}}, v_0, v_n, v_{n+2} \rangle; \quad {}_n\Phi_0^3({}_nA_3^{n-2}) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, \\ & v_{n-2}, v_0, v_1, v_0, \underline{v_2}, v_0, \dots, v_{n-3}, v_0, v_n, v_{n+2} \rangle. \end{aligned}$$
$$\begin{aligned} & \text{By Proposition 4.10 and Proposition 4.11, } {}_n\Phi_0^3(nA_3^i) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, \\ & v_0 \rangle \vee e_1^{n-1,2} \vee e_2^{n-1,2} \vee \cdots \vee e_{k_{n-1,2}}^{n-1,2} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n \rangle \\ & \vee \langle v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}}, \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \vee e_{k_{n-1,2}}^{n-1,2} \vee e_{k_{n-1,2}-1}^{n-1,2} \vee \cdots \vee e_1^{n-1,2} \\ & \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{j=i}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{j,2} \vee e_2^{j,2} \vee \cdots \vee e_{k_{j,2}}^{j,2} \\ & \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n, v_{n+2} \rangle \vee \langle v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}}, \\ & \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \vee e_{k_{j,2}}^{j,2} \vee e_{k_{j,2}-1}^{j,2} \vee \cdots \vee e_1^{j,2} \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, \\ & v_0 \rangle) \vee \langle v_0, v_{n-1}, v_0, v_n, v_{n+2}, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{j=1}^{i-1} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{j,2} \vee \\ & e_2^{j,2} \vee \cdots \vee e_{k_{j,2}}^{j,2} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n, v_{n+2} \rangle \vee \langle v_{n+2}, v_n, \\ & v_0, v_{n-3}, v_0, \underline{v_{n-4}}, \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \vee e_{k_{j,2}}^{j,2} \vee e_{k_{j,2}-1}^{j,2} \vee \cdots \vee e_1^{j,2} \vee \langle v_0, v_{n-1}, \\ & v_0, v_n, v_0, v_{n-1}, v_0 \rangle) \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{n,2} \vee e_2^{n,2} \vee \cdots \vee e_{k_{n,2}}^{n,2} \vee \langle v_0, v_{n-1}, \\ & v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n, v_{n+2} \rangle = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{n-1,2} \\ & \vee e_2^{n-1,2} \vee \cdots \vee e_{k_{n-1,2}}^{n-1,2} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n, v_0, v_{n-3}, v_0, \\ & \underline{v_{n-4}}, \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \vee e_{k_{n-1,2}}^{n-1,2} \vee e_{k_{n-1,2}-1}^{n-1,2} \vee \cdots \vee e_1^{n-1,2} \vee \langle v_0, v_{n-1}, v_0, \\ & v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{j=i}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{j,2} \vee e_2^{j,2} \vee \cdots \vee e_{k_{j,2}}^{j,2} \vee \langle v_0, v_{n-1}, v_0, \\ & v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n, v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}}, \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, v_{n-2}, v_0, \\ & v_{n-1}, v_0 \rangle \vee e_{k_{j,2}}^{j,2} \vee e_{k_{j,2}-1}^{j,2} \vee \cdots \vee e_1^{j,2} \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle) \vee \langle v_0, v_{n-1}, v_0, v_n, \\ & v_{n+2}, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{j=1}^{i-1} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{j,2} \vee e_2^{j,2} \vee \cdots \vee e_{k_{j,2}}^{j,2} \vee \langle v_0, \\ & v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, v_n, v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}}, \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, \\ & v_{n-2}, v_0, v_{n-1}, v_0 \rangle \vee e_{k_{j,2}}^{j,2} \vee e_{k_{j,2}-1}^{j,2} \vee \cdots \vee e_1^{j,2} \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee \langle v_0, v_{n-1}, \\ & v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{n,2} \vee e_2^{n,2} \vee \cdots \vee e_{k_{n,2}}^{n,2} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \\ & \underline{v_{n-3}}, \underline{v_0}, v_n, v_{n+2} \rangle \text{ for all } i \in \{1, \dots, n-3\}, \quad {}_n\Phi_0^3(nA_3^{n-1}) = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, \\ & v_0 \rangle \vee e_1^{n-1,2} \vee e_2^{n-1,2} \vee \cdots \vee e_{k_{n-1,2}}^{n-1,2} \vee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2}, \underline{v_0}, \dots, \underline{v_{n-3}}, \underline{v_0}, \\ & v_n \rangle \vee \langle v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}}, \underline{v_0}, \dots, \underline{v_1}, \underline{v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \vee e_{k_{n-1,2}}^{n-1,2} \vee e_{k_{n-1,2}-1}^{n-1,2} \vee \cdots \\ & \vee e_1^{n-1,2} \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \vee e_1^{n,2} \vee e_2^{n,2} \end{aligned}$$

$$\begin{aligned}
 & \bigvee \cdots \bigvee e_{k_{n,2}}^{n,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle = \langle v_0, v_{n-1}, v_0, \\
 & v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1,2} \bigvee e_2^{n-1,2} \bigvee \cdots \bigvee e_{k_{n-1,2}}^{n-1,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \\
 & \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1,2}}^{n-1,2} \bigvee e_{k_{n-1,2}-1}^{n-1,2} \bigvee \cdots \\
 & \bigvee e_1^{n-1,2} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n,2} \bigvee e_2^{n,2} \bigvee \\
 & \cdots \bigvee e_{k_{n,2}}^{n,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle, \overline{{}_n\Phi_0^3({}_nA_3^n)} = \langle v_0, \\
 & v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1,2} \bigvee e_2^{n-1,2} \bigvee \cdots \bigvee e_{k_{n-1,2}}^{n-1,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \\
 & \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n \rangle \bigvee \langle v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1,2}}^{n-1,2} \\
 & \bigvee e_{k_{n-1,2}-1}^{n-1,2} \bigvee \cdots \bigvee e_1^{n-1,2} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_{n+2}, v_n, v_0, \\
 & v_{n-1}, v_0 \rangle \bigvee_{i=i}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{i,2} \bigvee e_2^{i,2} \bigvee \cdots \bigvee e_{k_{i,2}}^{i,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, \\
 & v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle \bigvee \langle v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, \\
 & v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i,2}}^{i,2} \bigvee e_{k_{i,2}-1}^{i,2} \bigvee \cdots \bigvee e_1^{i,2} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle) \bigvee \langle v_0, v_{n-1}, \\
 & v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n,2} \bigvee e_2^{n,2} \bigvee \cdots \bigvee e_{k_{n,2}}^{n,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \\
 & \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1,2} \bigvee e_2^{n-1,2} \bigvee \cdots \bigvee e_{k_{n-1,2}}^{n-1,2} \bigvee \langle v_0, \\
 & v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, \\
 & v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1,2}}^{n-1,2} \bigvee e_{k_{n-1,2}-1}^{n-1,2} \bigvee \cdots \bigvee e_1^{n-1,2} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, \\
 & v_0, v_n, v_{n+2}, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{i=1}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{i,2} \bigvee e_2^{i,2} \bigvee \cdots \bigvee e_{k_{i,2}}^{i,2} \bigvee \\
 & \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \\
 & \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i,2}}^{i,2} \bigvee e_{k_{i,2}-1}^{i,2} \bigvee \cdots \bigvee e_1^{i,2} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle) \bigvee \\
 & \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n,2} \bigvee e_2^{n,2} \bigvee \cdots \bigvee e_{k_{n,2}}^{n,2} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \\
 & \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle; \overline{{}_n\Phi_0^4({}_nA_4^{n-2})} = \overline{{}_n\Phi_0^3({}_nA_3^{n-1})}. \text{ Thus, the claim is true} \\
 & \text{for } m = 4.
 \end{aligned}$$

Let $m \geq 4$ and suppose the claim is true for all $\tilde{m} \in \{3, \dots, m\}$. By Proposition 4.10 and Proposition 4.11, $\overline{{}_n\Phi_0^m({}_nA_m^i)} = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1, m-1} \bigvee$
 $e_2^{n-1, m-1} \bigvee \cdots \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n \rangle \bigvee \langle v_n, v_0,$
 $v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee e_{k_{n-1, m-1}-1}^{n-1, m-1} \bigvee \cdots \bigvee e_1^{n-1, m-1}$
 $\bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{j=0}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{i+j \bmod n-2, m-1} \bigvee$
 $e_2^{i+j \bmod n-2, m-1} \bigvee \cdots \bigvee e_{k_{i+j \bmod n-2, m-1}}^{i+j \bmod n-2, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0},$
 $v_n, v_{n+2} \rangle \bigvee \langle v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i+j \bmod n-2, m-1}}^{i+j \bmod n-2, m-1}$

$$\begin{aligned}
 & \bigvee e_{k_{i+j} \bmod n-2, m-1}^{i+j \bmod n-2, m-1} \bigvee \dots \bigvee e_1^{i+j \bmod n-2, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, \\
 & v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n, m-1} \bigvee e_2^{n, m-1} \bigvee \dots \bigvee e_{k_{n, m-1}}^{n, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \\
 & \dots, v_{n-3}, v_0, v_n, v_{n+2} \rangle = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1, m-1} \bigvee e_2^{n-1, m-1} \bigvee \dots \bigvee \\
 & e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \\
 & \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee e_{k_{n-1, m-1}-1}^{n-1, m-1} \bigvee \dots \bigvee e_1^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, \\
 & v_0, v_{n-1}, v_0 \rangle \bigvee_{j=0}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{i+j \bmod n-2, m-1} \bigvee e_2^{i+j \bmod n-2, m-1} \bigvee \dots \\
 & \bigvee e_{k_{i+j} \bmod n-2, m-1}^{i+j \bmod n-2, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{n-3}, \\
 & v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i+j} \bmod n-2, m-1}^{i+j \bmod n-2, m-1} \bigvee e_{k_{i+j} \bmod n-2, m-1-1}^{i+j \bmod n-2, m-1} \bigvee \dots \bigvee \\
 & e_1^{i+j \bmod n-2, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n, m-1} \\
 & \bigvee e_2^{n, m-1} \bigvee \dots \bigvee e_{k_{n, m-1}}^{n, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle \text{ for} \\
 & \text{all } i \in \{1, \dots, n-3\}, \quad \overline{n\Phi_0^m(nA_m^{n-1})} = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1, m-1} \bigvee \\
 & e_2^{n-1, m-1} \bigvee \dots \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n \rangle \bigvee \langle v_n, v_0, \\
 & v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee e_{k_{n-1, m-1}-1}^{n-1, m-1} \bigvee \dots \bigvee e_1^{n-1, m-1} \\
 & \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n, m-1} \bigvee e_2^{n, m-1} \bigvee \dots \\
 & \bigvee e_{k_{n, m-1}}^{n, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle = \langle v_0, v_{n-1}, v_0, v_n, \\
 & v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1, m-1} \bigvee e_2^{n-1, m-1} \bigvee \dots \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \\
 & \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee e_{k_{n-1, m-1}-1}^{n-1, m-1} \\
 & \bigvee \dots \bigvee e_1^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n, m-1} \\
 & \bigvee e_2^{n, m-1} \bigvee \dots \bigvee e_{k_{n, m-1}}^{n, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle, \\
 & \overline{n\Phi_0^m(nA_m^n)} = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1, m-1} \bigvee e_2^{n-1, m-1} \bigvee \dots \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee \\
 & \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n \rangle \bigvee \langle v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0} \\
 & , v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1, m-1}}^{n-1, m-1} \bigvee e_{k_{n-1, m-1}-1}^{n-1, m-1} \bigvee \dots \bigvee e_1^{n-1, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, \\
 & v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-2, m-1} \bigvee e_2^{n-2, m-1} \bigvee \dots \bigvee e_{k_{n-2, m-1}}^{n-2, m-1} \bigvee \langle v_0, v_{n-1}, \\
 & v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle \bigvee \langle v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \\
 & \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-2, m-1}}^{n-2, m-1} \bigvee e_{k_{n-2, m-1}-1}^{n-2, m-1} \bigvee \dots \bigvee e_1^{n-2, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, \\
 & v_0, v_{n-1}, v_0 \rangle \bigvee_{i=i}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{i, m-1} \bigvee e_2^{i, m-1} \bigvee \dots \bigvee e_{k_{i, m-1}}^{i, m-1} \bigvee \langle v_0, \\
 & v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle \bigvee \langle v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \\
 & \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i, m-1}}^{i, m-1} \bigvee e_{k_{i, m-1}-1}^{i, m-1} \bigvee \dots \bigvee e_1^{i, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, \\
 & \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i, m-1}}^{i, m-1} \bigvee e_{k_{i, m-1}-1}^{i, m-1} \bigvee \dots \bigvee e_1^{i, m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1},
 \end{aligned}$$

$v_0\rangle) \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n,m-1} \bigvee e_2^{n,m-1} \bigvee \cdots \bigvee e_{k_{n,m-1}}^{n,m-1} \bigvee \langle v_0, v_{n-1}, v_0,$
 $v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle = \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-1,m-1} \bigvee$
 $e_2^{n-1,m-1} \bigvee \cdots \bigvee e_{k_{n-1,m-1}}^{n-1,m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3},$
 $v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-1,m-1}}^{n-1,m-1} \bigvee e_{k_{n-1,m-1}-1}^{n-1,m-1} \bigvee \cdots \bigvee e_1^{n-1,m-1} \bigvee \langle v_0,$
 $v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n-2,m-1} \bigvee e_2^{n-2,m-1} \bigvee \cdots \bigvee$
 $e_{k_{n-2,m-1}}^{n-2,m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4},$
 $\underline{v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{n-2,m-1}}^{n-2,m-1} \bigvee e_{k_{n-2,m-1}-1}^{n-2,m-1} \bigvee \cdots \bigvee e_1^{n-2,m-1} \bigvee \langle v_0, v_{n-1},$
 $v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee_{i=1}^{n-3} (\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{i,m-1} \bigvee e_2^{i,m-1} \bigvee \cdots \bigvee e_{k_{i,m-1}}^{i,m-1} \bigvee \langle v_0,$
 $v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0},$
 $v_{n-2}, v_0, v_{n-1}, v_0 \rangle \bigvee e_{k_{i,m-1}}^{i,m-1} \bigvee e_{k_{i,m-1}-1}^{i,m-1} \bigvee \cdots \bigvee e_1^{i,m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle) \bigvee$
 $\langle v_0, v_{n-1}, v_0, v_n, v_0, v_{n-1}, v_0 \rangle \bigvee e_1^{n,m-1} \bigvee e_2^{n,m-1} \bigvee \cdots \bigvee e_{k_{n,m-1}}^{n,m-1} \bigvee \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0,$
 $v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle; \quad \overline{{}_n\Phi_0^{m+1}}({}_nA_{m+1}^{n-2}) = \overline{{}_n\Phi_0^m}({}_nA_m^{n-1}),$ and so the claim
 holds by induction. $\square$

4.3 Factoring

In this section, the complexity of ${}_nK$ is addressed by resolving whether or not it is simpler (Proposition 4.20). Were it to be, in keeping with the strategy by Minc, a contradiction to a factoring such as given by Theorem 3.3 is obtained. In line with what was developed in [15], a certain factoring allowing for control of the complexity of the dual system is deduced, through properties of the bonding map, when given a factoring in general. In particular, a factoring such that the dual of the simpler object through which the bonding map is being factored remains simpler, deduced, here, from the more general factoring in which branch point of the simpler object goes to branch point. For ${}_nK$, the symmetry of the bonding maps described in the previous section (Proposition 4.12) facilitates this with Propositions 4.16 and 4.17. The argument of [15] is then further adapted, with the conditions of Minc's program being satisfied as established in the previous section, to allow for a "shortening" and "lifting" of such a factoring after passing to the dual (Proposition 4.18), showing that a factoring cannot occur (Proposition 4.19), provided that certain induction

conditions are preserved and that the defining bonding map $({}_n\phi)$ cannot be factored.

For Propositions 4.13 through 4.17, let $m \geq 2$, $\tilde{n} \in \{1, \dots, n-1\}$, and T be a simple- $\tilde{n}$ -od. Let $S^1, S^2, \dots, S^{\tilde{n}}$ be $\tilde{n}$ arcs so that $\bigcup_{i=1}^{\tilde{n}} S^i = T$ and $S^i \cap S^j = \{s_0\}$ for each distinct $i, j \in \{1, \dots, \tilde{n}\}$, and so s_0 is an endpoint of S^i for each $i \in \{1, \dots, \tilde{n}\}$. Define T as a graph with subgraph $S^i = \langle s_0, s_1^i, \dots, s_{k_i}^i \rangle$ for some $k_i \in \{1, 2, \dots\}$ for each $i \in \{1, \dots, \tilde{n}\}$. Suppose $\alpha : {}_nX_m \longrightarrow T$ and $\beta : T \longrightarrow {}_nX_0$ are simplicial maps so that α is surjective, $\beta \circ \alpha = {}_n\Phi_0^m$, and $\beta(s_0) = v_0$. Let ${}_nA_m^i = \langle x_0^i, x_1^i, \dots, x_{\ell_i}^i \rangle$ for some $\ell_i \in \{1, 2, \dots\}$ for each $i \in \{1, \dots, n\}$ where $x_0^i = v_0$ for all $i \in \{1, \dots, n\}$.

In the case of the existence of distinct $x, y, z \in V({}_nX_m)$ so that x and y are adjacent, y and z are adjacent, $\alpha(y) = s_0$, ${}_n\Phi_0^m(z) \neq v_{n-1}$, and ${}_n\Phi_0^m(x) = v_n$, Propositions 4.13 through 4.15 show that for all vertices of ${}_nX_m$ mapped by α to s_0 a certain uniqueness applies in describing how they are “locally” mapped by ${}_n\Phi_0^m$. This uniqueness ensures certain instances, in which the above case holds, do not prevent the “controllable” factoring arranged in Proposition 4.16 from being established as well-defined.

Proposition 4.13. *Let $i \in \{1, \dots, n\}$, $r \in \{0, \dots, \ell_i\}$, and $j \in \{1, \dots, n-2\}$ so that $\overline{{}_n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $\overline{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+t}^i \rangle) = \langle s_{t-1}^i, s_{t-2}^i, \dots, s_1^i, s_0 \rangle$ for some even $t \in \{2, \dots, 2(n-2)\}$ and for some $\hat{i} \in \{1, \dots, \tilde{n}\}$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$. Then,*

i) there exists $r' \in \{0, \dots, \ell_i\}$ so that $r' < r$, $\overline{{}_n\Phi_0^m}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $\overline{\alpha}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+t'}^i \rangle) = \langle s_{t'-1}^i, s_{t'-2}^i, \dots, s_1^i, s_0 \rangle$ for some even $t' \in \{2, \dots, 2(n-2)\}$ and for some $\hat{i}' \in \{1, \dots, \tilde{n}\}$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, or

ii) there exists $r' \in \{0, \dots, \ell_i\}$ so that $r' < r$, $\overline{{}_n\Phi_0^m}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+4(n-2)+5}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{p^j(n-2)}, v_0, \underline{v_{p^j(n-3)}, v_0}, \dots, \underline{v_{p^j(1)}, v_0}, v_{n-1} \rangle$ (or $\overline{{}_n\Phi_0^m}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+4(n-2)+3}^i \rangle) = \langle v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1} \rangle$), $\overline{\alpha}(\langle x_{r'+4(n-2)+5-(t+1)}^i, x_{r'+4(n-2)+5-t}^i, \dots, x_{r'+4(n-2)+5}^i \rangle) = \langle s_0, s_1^i, \dots, s_{t+1}^i \rangle$ (or $\overline{\alpha}(\langle x_{r'+4(n-2)+3-(t+1)}^i, x_{r'+4(n-2)+3-t}^i, \dots, x_{r'+4(n-2)+3}^i \rangle) = \langle s_0, s_1^i, \dots, s_{t+1}^i \rangle$).

$\rangle) = \langle s_0, s_1^i, \dots, s_{t+1}^i \rangle$, and $\overline{\alpha}(\langle x_0^i, x_1^i, \dots, x_{r'+2(n-2)+1}^i \rangle) \subseteq S^{\tilde{i}'}$ where $\tilde{i}' \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{r'+1}^i) \in S^{\tilde{i}'}$ or

$$iii) \overline{\alpha}(\langle x_0^i, x_1^i, \dots, x_r^i \rangle) \subseteq S^{\tilde{i}}.$$

Proof. Suppose $\overline{\alpha}(\langle x_0^i, x_1^i, \dots, x_r^i \rangle) \not\subseteq S^{\tilde{i}}$. By Proposition 4.12, $\overline{n\Phi_0^m}(\langle x_{r-1}^i, x_r^i \rangle) = \langle v_{n-1}, v_0 \rangle$. If $t = 2$, then $|\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\}| = \tilde{n}$. Hence, $\overline{\alpha}(\langle x_{r-1}^i, x_r^i \rangle) = \langle s_{t+1}^{\tilde{i}}, s_t^{\tilde{i}} \rangle$. Then, by Proposition 4.12, there exists $\tilde{r}' < r$ so that $\overline{n\Phi_0^m}(\langle x_{\tilde{r}'+1}^i, x_{\tilde{r}'+2}^i, \dots, x_{\tilde{r}'+4(n-2)+5}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{p^j(n-2)}, v_0, \underline{v_{p^j(n-3)}, v_0}, \dots, \underline{v_{p^j(1)}, v_0}, v_{n-1} \rangle$ (or $\overline{n\Phi_0^m}(\langle x_{\tilde{r}'+1}^i, x_{\tilde{r}'+2}^i, \dots, x_{\tilde{r}'+4(n-2)+3}^i \rangle) = \langle v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1} \rangle$) and $\overline{\alpha}(\langle x_{\tilde{r}'+4(n-2)+5-(t+1)}^i, x_{\tilde{r}'+4(n-2)+5-t}^i, \dots, x_{\tilde{r}'+4(n-2)+5}^i \rangle) = \langle s_0, s_1^{\tilde{i}}, \dots, s_{t+1}^{\tilde{i}} \rangle$ (or $\overline{\alpha}(\langle x_{\tilde{r}'+4(n-2)+3-(t+1)}^i, x_{\tilde{r}'+4(n-2)+3-t}^i, \dots, x_{\tilde{r}'+4(n-2)+3}^i \rangle) = \langle s_0, s_1^{\tilde{i}}, \dots, s_{t+1}^{\tilde{i}} \rangle$).

Let $\tilde{i}' \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{\tilde{r}'+1}^i) \in S^{\tilde{i}'}$.

Case 1: $\alpha(x_{\tilde{r}'+2(n-2)+1}^i) \in S^{\tilde{i}'}$.

Then, there exists an even $\tilde{t}' \in \{t, \dots, 2(n-2)\}$ so that $\alpha(x_{\tilde{r}'+4(n-2)+5-(\tilde{t}'+2)}^i) = s_1^{\tilde{i}'}$ (or $\alpha(x_{\tilde{r}'+4(n-2)+3-(\tilde{t}'+2)}^i) = s_1^{\tilde{i}'}$). If $\overline{\alpha}(\langle x_0^i, x_1^i, \dots, x_{\tilde{r}'}^i \rangle) \subseteq S^{\tilde{i}'}$, then conclusion (ii) holds with $r' = \tilde{r}'$ and $\tilde{i}' = \tilde{i}'$. Suppose otherwise. By Proposition 4.12 there exists $r' < \tilde{r}'$ so that $\overline{n\Phi_0^m}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+2(n-2)+1}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n \rangle$ and $\overline{\alpha}(\langle x_{r'+\tilde{t}'}^i, x_{r'+\tilde{t}'+1}^i \rangle) = \langle s_0, s_1^{\tilde{i}'} \rangle$. Let $\tilde{i}' \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{r'+1}^i) \in S^{\tilde{i}'}$. Then, there exists an even $t' \in \{2, \dots, \tilde{t}'\}$ so that $\overline{\alpha}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+t'}^i \rangle) = \langle s_{t'-1}^{\tilde{i}'}, s_{t'-2}^{\tilde{i}'}, \dots, s_1^{\tilde{i}'}, s_0 \rangle$ and $\{\alpha(x_{r'+\tilde{t}}^i) : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, \tilde{t}'\}\} \subseteq \{s_1^\delta : \delta \in \{1, \dots, \tilde{n}\}\}$, giving $\{v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, \tilde{t}'\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$. Since $\tilde{t}' \geq t$ and by hypothesis $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, then $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, and conclusion (i) is reached.

Case 2: $\alpha(x_{\tilde{r}'+2(n-2)+1}^i) \notin S^{\tilde{i}'}$.

Then, there exists an even $t' \in \{2, \dots, 2(n-2)\}$ so that $\overline{\alpha}(\langle x_{\tilde{r}'+1}^i, x_{\tilde{r}'+2}^i, \dots, x_{\tilde{r}'+t'}^i \rangle) = \langle s_{t'-1}^{\tilde{i}'}, s_{t'-2}^{\tilde{i}'}, \dots, s_1^{\tilde{i}'} s_0 \rangle$. Let $\tilde{i}' \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{\tilde{r}'+2(n-2)+1}^i) \in S^{\tilde{i}'}$. Then, there exists an even $\tilde{t}' \in \{\max\{t, t'\}, \dots, 2(n-2)\}$ so that $\overline{\alpha}(\langle x_{\tilde{r}'+\tilde{t}'}^i, x_{\tilde{r}'+\tilde{t}'+1}^i \rangle) = \langle s_0, s_1^{\tilde{i}'} \rangle$ and $\{\alpha(x_{\tilde{r}'+\tilde{t}}^i) : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, \tilde{t}'\}\} \subseteq \{s_1^\delta : \delta \in \{1, \dots, \tilde{n}\}\}$, giving

$\{v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, \tilde{t}'\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$. Since $\tilde{t}' \geq t$ and by hypothesis $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, then $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, and conclusion (i) holds with $r' = \tilde{r}'$ and $\hat{i}' = \tilde{i}'$. $\square$

Proposition 4.14. *Let $x \in V(nX_m)$ so that $\alpha(x) = s_0$. Suppose there exist $i \in \{1, \dots, n\}$, $r \in \{0, \dots, \ell_i\}$, and $j \in \{1, \dots, n-2\}$ so that $\overline{n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $\overline{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+t}^i \rangle) = \langle s_{t-1}^i, s_{t-2}^i, \dots, s_1^i, s_0 \rangle$ for some even $t \in \{2, \dots, 2(n-2)\}$ and for some $i \in \{1, \dots, \tilde{n}\}$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$. Then, there exist $i \in \{1, \dots, n\}$, $\hat{r} \in \{0, \dots, \ell_i\}$, and an even $\hat{t} \in \{2, \dots, 2(n-2)\}$ so that*

- i) $\overline{n\Phi_0^m}(\langle x_{\hat{r}+1}^i, x_{\hat{r}+2}^i, \dots, x_{\hat{r}+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $x = x_{\hat{r}+\hat{t}}^i$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{\hat{t}, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, or
- ii) $\overline{n\Phi_0^m}(\langle x_{\hat{r}-1}^i, x_{\hat{r}-2}^i, \dots, x_{\hat{r}-2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $x = x_{\hat{r}-\hat{t}}^i$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{\hat{t}, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$.

Proof. By Proposition 4.13, there exists $r' \in \{0, \dots, \ell_i\}$ so that $r' \leq r$, $\overline{n\Phi_0^m}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $\overline{\alpha}(\langle x_{r'+1}^i, x_{r'+2}^i, \dots, x_{r'+t'}^i \rangle) = \langle s_{t'-1}^i, s_{t'-2}^i, \dots, s_1^i, s_0 \rangle$ for some even $t' \in \{2, \dots, 2(n-2)\}$ and for some $\hat{i}' \in \{1, \dots, \tilde{n}\}$, $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t', \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, and a) $\overline{\alpha}(\langle x_0^i, x_1^i, \dots, x_{r'}^i \rangle) \subseteq S^{\hat{i}'}$ or b) there exists $\tilde{r}' \in \{0, \dots, \ell_i\}$ so that $\tilde{r}' < r'$, $\overline{n\Phi_0^m}(\langle x_{\tilde{r}'+1}^i, x_{\tilde{r}'+2}^i, \dots, x_{\tilde{r}'+4(n-2)+5}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0}, v_n, v_{n+2}, v_n, v_0, v_{p^j(n-2)}, v_0, \underline{v_{p^j(n-3)}, v_0}, \dots, \underline{v_{p^j(1)}, v_0}, v_{n-1} \rangle$ (or $\overline{n\Phi_0^m}(\langle x_{\tilde{r}'+1}^i, x_{\tilde{r}'+2}^i, \dots, x_{\tilde{r}'+4(n-2)+3}^i \rangle) = \langle v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_0, v_{n-3}, v_0, \underline{v_{n-4}, v_0}, \dots, \underline{v_1, v_0}, v_{n-2}, v_0, v_{n-1} \rangle$), $\overline{\alpha}(\langle x_{\tilde{r}'+4(n-2)+5-(t'+1)}^i, x_{\tilde{r}'+4(n-2)+5-t'}^i, \dots, x_{\tilde{r}'+4(n-2)+5}^i \rangle) = \langle s_0, s_1^{\hat{i}'}, \dots, s_{t'+1}^{\hat{i}'} \rangle$ (or $\overline{\alpha}(\langle x_{\tilde{r}'+4(n-2)+3-(t'+1)}^i, x_{\tilde{r}'+4(n-2)+3-t'}^i, \dots, x_{\tilde{r}'+4(n-2)+3}^i \rangle) = \langle s_0, s_1^{\hat{i}'}, \dots, s_{t'+1}^{\hat{i}'} \rangle$), and $\overline{\alpha}(\langle x_0^i, x_1^i, \dots, x_{\tilde{r}'+2(n-2)+1}^i \rangle) \subseteq S^{\tilde{i}'}$ where $\tilde{i}' \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{\tilde{r}'+1}^i) \in S^{\tilde{i}'}$.

Let $\check{i} \in \{1, \dots, n\}$ and $\check{r} \in \{0, \dots, \ell_i\}$ so that $\alpha^{-1}(\{s_0\}) \cap \langle x_0^{\check{i}}, x_1^{\check{i}}, \dots, x_{\check{r}}^{\check{i}} \rangle = \{x_{\check{r}}^{\check{i}}\}$. If $\check{r}$ exists and (a) holds, then by the hypothesis and Proposition 4.12, conclusion (i) of the claim holds for $x_{\check{r}}^{\check{i}}$ with $i = \check{i}$, $\hat{r} = \check{r} - t'$, and $\hat{t} = t'$. Suppose $\check{r}$ exists and (b)

holds, and let $\tilde{t}' \in \{2(n-2)+4, \dots, 4(n-2)+4-t'\}$ (or $\tilde{t}' \in \{2(n-2)+2, \dots, 4(n-2)+2-t'\}$) so that $\alpha^{-1}(\{s_0\}) \cap \langle x_0^i, x_1^i, \dots, x_{\tilde{r}'+\tilde{t}'}^i \rangle = \{x_{\tilde{r}'+\tilde{t}'}^i\}$. Then by the hypothesis and Proposition 4.12, conclusion (ii) of the claim holds for $x_{\tilde{r}}^{\check{i}}$ with $i = \check{i}$, $\check{r} = \check{r} + 4(n-2) + 4 - \tilde{t}'$ (or $\check{r} = \check{r} + 4(n-2) + 2 - \tilde{t}'$), and $\check{t} = 4(n-2) + 4 - \tilde{t}'$ (or $\check{t} = 4(n-2) + 2 - \tilde{t}'$).

Suppose $\check{r}$ exists and let $\check{r}_\theta \in \{\check{r} + 1, \dots, \ell_{\check{i}}\}$ so that the claim holds for all $x \in \{x_{\check{r}}^{\check{i}} : \check{r} \in \{0, \dots, \check{r}_\theta - 1\} \text{ and } \alpha(x_{\check{r}}^{\check{i}}) = s_0\}$ and $\alpha(x_{\check{r}_\theta}^{\check{i}}) = s_0$. By Proposition 4.12, $\alpha(x_{\check{r}_\theta-1}^{\check{i}}) = s_1^{\hat{i}_-}$ for some $\hat{i}_- \in \{1, \dots, \tilde{n}\}$ and $\alpha(x_{\check{r}_\theta+1}^{\check{i}}) = s_1^{\hat{i}_+}$ for some $\hat{i}_+ \in \{1, \dots, \tilde{n}\}$.

Case i: $\alpha(x_{\check{r}_\theta-2}^{\check{i}}) = s_0$.

By the induction hypothesis, there exist $\check{r}_0 \in \{0, \dots, \ell_{\check{i}}\}$ and an even $\check{t}_0 \in \{2, \dots, 2(n-2)\}$ so that

1) $\overline{{}_n\Phi_0^m}(\langle x_{\check{r}_0+1}^{\check{i}}, x_{\check{r}_0+2}^{\check{i}}, \dots, x_{\check{r}_0+2(n-2)}^{\check{i}} \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $\check{r}_\theta - 2 = \check{r}_0 + \check{t}_0$, and $\{v_n, v_{p^j(\frac{\check{t}}{2})} : \check{t} \text{ is even and } \check{t} \in \{\check{t}_0, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, or

2) $\overline{{}_n\Phi_0^m}(\langle x_{\check{r}_0-1}^{\check{i}}, x_{\check{r}_0-2}^{\check{i}}, \dots, x_{\check{r}_0-2(n-2)}^{\check{i}} \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$, $\check{r}_\theta - 2 = \check{r}_0 - \check{t}_0$, and $\{v_n, v_{p^j(\frac{\check{t}}{2})} : \check{t} \text{ is even and } \check{t} \in \{\check{t}_0, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$.

Case i.a: Suppose (1) holds.

If $\check{t}_0 \in \{2, \dots, 2(n-2) - 2\}$, then conclusion (i) of the claim holds for $x_{\check{r}_\theta}^{\check{i}}$ with $i = \check{i}$, $\check{r} = \check{r}_0$, and $\check{t} = \check{t}_0 + 2$. If $\check{t}_0 = 2(n-2)$, then by Proposition 4.12, conclusion (ii) of the claim holds for $x_{\check{r}_\theta}^{\check{i}}$ with $i = \check{i}$, $\check{r} = \check{r}_0 + 4(n-2) + 2$, and $\check{t} = 2(n-2)$.

Case i.b: Suppose (2) holds.

If $\check{t}_0 \in \{4, \dots, 2(n-2)\}$, then conclusion (ii) of the claim holds for $x_{\check{r}_\theta}^{\check{i}}$ with $i = \check{i}$, $\check{r} = \check{r}_0$, and $\check{t} = \check{t}_0 - 2$. If $\check{t}_0 = 2$, then $|\{v_n, v_{p^j(\frac{\check{t}}{2})} : \check{t} \text{ is even and } \check{t} \in \{\check{t}_0, \dots, 2(n-2)\}\}| = \tilde{n}$, but $\beta(s_1^{\hat{i}_+}) = v_{n-1}$, since by Proposition 4.12 ${}_n\Phi_0^m(x_{\check{r}_\theta+1}^{\check{i}}) = v_{n-1}$, giving a contradiction.

Case ii: $\alpha(x_{\check{r}_\theta-2}^{\check{i}}) \neq s_0$.

Then, $\alpha(x_{\check{r}_\theta-2}^{\check{i}}) = s_2^{\hat{i}_-}$, and by the induction hypothesis, there exist $\check{r}_0 \in \{0, \dots, \ell_{\check{i}}\}$ and an even $\check{t}_0 \in \{2, \dots, 2(n-2)\}$ so that

1) $\acute{r}_0 + \acute{t}_0 < \check{r}_\theta$, $\overline{{}_n\Phi_0^m}(\langle x_{\acute{r}_0+1}^{\check{i}}, x_{\acute{r}_0+2}^{\check{i}}, \dots, x_{\acute{r}_0+2(n-2)}^{\check{i}} \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, v_{p^j(n-2)}, v_0 \rangle$, $\overline{\alpha}(\langle x_{\acute{r}_0+\acute{t}_0}^{\check{i}}, x_{\acute{r}_0+\acute{t}_0+1}^{\check{i}}, x_{\acute{r}_0+\acute{t}_0+2}^{\check{i}} \rangle) = \langle s_0, s_1^{\hat{i}_-}, s_2^{\hat{i}_-} \rangle$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{\acute{t}_0, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, or

2) $\acute{r}_0 - \acute{t}_0 < \check{r}_\theta$, $\overline{{}_n\Phi_0^m}(\langle x_{\acute{r}_0-1}^{\check{i}}, x_{\acute{r}_0-2}^{\check{i}}, \dots, x_{\acute{r}_0-2(n-2)}^{\check{i}} \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, v_{p^j(n-2)}, v_0 \rangle$, $\overline{\alpha}(\langle x_{\acute{r}_0-\acute{t}_0}^{\check{i}}, x_{\acute{r}_0-\acute{t}_0+1}^{\check{i}}, x_{\acute{r}_0-\acute{t}_0+2}^{\check{i}} \rangle) = \langle s_0, s_1^{\hat{i}_-}, s_2^{\hat{i}_-} \rangle$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{\acute{t}_0, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$.

Case ii.a: Suppose (1) holds.

By Proposition 4.12, conclusion (ii) of the claim holds for $x_{\check{r}_\theta}^{\check{i}}$ with $i = \check{i}$, $\acute{r} = \check{r}_\theta + \acute{t}_0$, and $\acute{t} = \acute{t}_0$.

Case ii.b: Suppose (2) holds.

By Proposition 4.12, conclusion (i) of the claim holds for $x_{\check{r}_\theta}^{\check{i}}$ with $i = \check{i}$, $\acute{r} = \check{r}_\theta - \acute{t}_0$, and $\acute{t} = \acute{t}_0$.

Thus, the claim follows by induction. $\square$

Proposition 4.15. *Let distinct $x, y, z \in V({}_nX_m)$ be so that x and y are adjacent, y and z are adjacent, $\alpha(y) = s_0$, ${}_n\Phi_0^m(z) \neq v_{n-1}$, and ${}_n\Phi_0^m(x) = v_n$. Then,*

i) *there exist $i \in \{1, \dots, n\}$, $r \in \{0, \dots, \ell_i\}$, and $j \in \{1, \dots, n-2\}$ so that $\overline{{}_n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, v_{p^j(n-2)}, v_0 \rangle$, $\overline{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+t}^i \rangle) = \langle s_{t-1}^{\hat{i}_-}, s_{t-2}^{\hat{i}_-}, \dots, s_1^{\hat{i}_-}, s_0 \rangle$ for some even $t \in \{2, \dots, 2(n-2)\}$ and for some $\hat{i} \in \{1, \dots, \tilde{n}\}$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$, or*

ii) *there exists $j \in \{1, \dots, n-2\}$ so that if $\tilde{x} \in V({}_nX_m)$ and $\alpha(\tilde{x}) = s_0$, then there exist $i \in \{1, \dots, n\}$ and $r \in \{0, \dots, \ell_i\}$ so that $\overline{{}_n\Phi_0^m}(\langle x_{r-1}^i, x_{r-2}^i, \dots, x_{r-2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, v_{p^j(n-2)}, v_0 \rangle$, $\tilde{x} = x_{r-t}^i$ for some even $t \in \{2, \dots, 2(n-2)\}$, and $\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t, \dots, 2(n-2)\}\} \subseteq \{\beta(s_1^\delta) : \delta \in \{1, \dots, \tilde{n}\}\}$.*

Proof. By Proposition 4.12, there exist $i' \in \{1, \dots, n\}$ and $r' \in \{0, \dots, \ell_{i'}\}$ so that for some $j \in \{1, \dots, n-2\}$,

a) $\overline{{}_n\Phi_0^m}(\langle x_{r'+1}^{i'}, x_{r'+2}^{i'}, \dots, x_{r'+2(n-2)}^{i'} \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, v_{p^j(n-2)}, v_0 \rangle$ and $\langle x, y, z \rangle = \langle x_{r'+2(n-2)+1}^{i'}, x_{r'+2(n-2)}^{i'}, x_{r'+2(n-2)-1}^{i'} \rangle$, or

b) $\overline{{}_n\Phi_0^m}(\langle x_{r'-1}^{i'}, x_{r'-2}^{i'}, \dots, x_{r'-2(n-2)}^{i'} \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$ and $\langle x, y, z \rangle = \langle x_{r'-2(n-2)-1}^{i'}, x_{r'-2(n-2)}^{i'}, x_{r'-2(n-2)+1}^{i'} \rangle$.

If (a) holds, then there exist an even $t \in \{2, \dots, 2(n-2)\}$ and $\hat{i} \in \{1, \dots, \tilde{n}\}$ so that the conditions of conclusion (i) are satisfied with $r = r'$ and $i = i'$. Suppose (b) is the case, and let $\hat{i}' \in \{1, \dots, \tilde{n}\}$ be so that $\alpha(x) = s_1^{\hat{i}'}$. If $\alpha(\langle x_0^{i'}, x_1^{i'}, \dots, x_{r'-2(n-2)-1}^{i'} \rangle) \not\subseteq S^{\hat{i}'}$, then by Proposition 4.12, there exist $r \in \{0, \dots, r' - 4(n-2) - 4\}$ (or $r \in \{0, \dots, r' - 4(n-2) - 2\}$), an even $t \in \{2, \dots, 2(n-2)\}$, and $\hat{i} \in \{1, \dots, \tilde{n}\}$ so that the conditions of conclusion (i) are satisfied with $i = i'$. Suppose $\alpha(\langle x_0^{i'}, x_1^{i'}, \dots, x_{r'-2(n-2)-1}^{i'} \rangle) \subseteq S^{\hat{i}'}$. By Proposition 4.12, if there exist $i \in \{1, \dots, n\}$ and $r_i \in \{0, \dots, \ell_i\}$ so that $\alpha(x_{r_i}^i) = s_0$ and $\alpha(\langle x_0^i, x_1^i, \dots, x_{r_i}^i \rangle) \subseteq S^{\hat{i}'}$, then $\alpha^{-1}(\{s_0\}) \cap \langle x_0^i, x_1^i, \dots, x_{r_i}^i \rangle = \{x_{r_i}^i\}$ and $\overline{{}_n\Phi_0^m}(\langle x_{\tilde{r}+1}^i, x_{\tilde{r}+2}^i, \dots, x_{\tilde{r}+2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$ where $\tilde{r} = r_i - 2(n-2) - 4$ (or $\tilde{r} = r_i - 2(n-2) - 2$). Suppose r_i exists for some $i \in \{1, \dots, n\}$. Let $\tilde{r}' = \tilde{r} + 4(n-2) + 4$ (or $\tilde{r}' = \tilde{r} + 4(n-2) + 2$). By Proposition 4.12, $\overline{{}_n\Phi_0^m}(\langle x_{\tilde{r}'-1}^i, x_{\tilde{r}'-2}^i, \dots, x_{\tilde{r}'-2(n-2)}^i \rangle) = \langle v_{p^j(1)}, v_0, \underline{v_{p^j(2)}, v_0}, \dots, \underline{v_{p^j(n-2)}, v_0} \rangle$. Let $\hat{i} \in \{1, \dots, \tilde{n}\} \setminus \{\hat{i}'\}$ so that $\overline{\alpha}(\langle x_{\tilde{r}'-1}^i, x_{\tilde{r}'-2}^i, \dots, x_{\tilde{r}'-t_i}^i \rangle) = \langle s_{t_i-1}^{\hat{i}}, s_{t_i-2}^{\hat{i}}, \dots, s_1^{\hat{i}}, s_0 \rangle$ for some even $t_i \in \{2, \dots, 2(n-2)\}$. By Proposition 4.12, $\overline{{}_n\Phi_0^m}(\langle x_{\tilde{r}'}^i, x_{\tilde{r}'+1}^i \rangle) = \langle v_0, v_{n-1} \rangle$. If $t_i = 2$, then $|\{v_n, v_{p^j(\frac{\tilde{t}}{2})} : \tilde{t} \text{ is even and } \tilde{t} \in \{t_i, \dots, 2(n-2)\}\}| = \tilde{n}$. Hence, $\overline{\alpha}(\langle x_{\tilde{r}'}^i, x_{\tilde{r}'+1}^i \rangle) = \langle s_{t_i}^{\hat{i}}, s_{t_i+1}^{\hat{i}} \rangle$. If there exists $\tilde{r}'_i \in \{\tilde{r}' + 1, \dots, \ell_i\}$ so that $\langle x_{\tilde{r}'}^i, x_{\tilde{r}'+1}^i, \dots, x_{\tilde{r}'_i}^i \rangle \subseteq S^{\hat{i}}$ and $\alpha(x_{\tilde{r}'_i}^i) = s_0$, then by Proposition 4.12, the conditions of conclusion (i) are satisfied with $r = \tilde{r}'_i - t_i$ and $t = t_i$. Suppose $\tilde{r}'_i$ does not exist for each $i \in \{1, \dots, n\}$, and let $\tilde{x} \in V({}_nX_m)$ so that $\alpha(\tilde{x}) = s_0$. Then, there exist $i \in \{1, \dots, n\}$ and some even $t \in \{t_i, \dots, 2(n-2)\}$ so that the conditions of conclusion (ii) are satisfied with $r = r_i + 2(n-2)$. $\square$

Proposition 4.16. *If $m, \tilde{n}, T, \alpha$, and β exist, then for some $\tilde{n}_0 \in \{1, \dots, \tilde{n}\}$, there exist a simple- $\tilde{n}_0$ -od $\tilde{T}$, $\tilde{n}_0$ arcs $\tilde{S}^1, \tilde{S}^2, \dots, \tilde{S}^{\tilde{n}_0}$ so that $\bigcup_{i=1}^{\tilde{n}_0} \tilde{S}^i = \tilde{T}$, $\tilde{S}^i \cap \tilde{S}^j = \{\tilde{s}_0\}$ for each distinct $i, j \in \{1, \dots, \tilde{n}_0\}$, so $\tilde{s}_0$ is an endpoint of $\tilde{S}^i$ for each $i \in \{1, \dots, \tilde{n}_0\}$, and $\tilde{S}^i = \langle \tilde{s}_0, \tilde{s}_1^i, \dots, \tilde{s}_{\tilde{k}_i}^i \rangle$ for some $\tilde{k}_i \in \{1, 2, \dots\}$ for each $i \in \{1, \dots, \tilde{n}_0\}$, and simplicial maps $\tilde{\alpha} : {}_nX_m \longrightarrow \tilde{T}$ and $\tilde{\beta} : \tilde{T} \longrightarrow {}_nX_0$ so that $\tilde{\beta} \circ \tilde{\alpha} = {}_n\Phi_0^m$, $\tilde{\beta}(\tilde{s}_0) = v_0$, and if*

distinct $x, y, z \in V(nX_m)$ are such that x and y are adjacent, y and z are adjacent, and $\tilde{\alpha}(y) = \tilde{s}_0$, then ${}_n\Phi_0^m(x) = v_{n-1}$ or ${}_n\Phi_0^m(z) = v_{n-1}$.

Proof. Let $D = \{\delta \in \{1, \dots, \tilde{n}\} : \text{there exist distinct } x, y, z \in V(nX_m) \text{ so that } \langle x, y, z \rangle \subseteq {}_nX_m, \alpha(y) = s_0, {}_n\Phi_0^m(x) \neq v_{n-1}, {}_n\Phi_0^m(z) \neq v_{n-1}, \text{ and } \alpha(x) = s_1^\delta\}$, $D_0 = \{\delta \in D : s_2^\delta \text{ does not exist and } \beta(s_1^\delta) \neq v_n\}$, $D_1 = \{\delta \in D : s_2^\delta \text{ exists and } \bar{\beta}(\langle s_1^\delta, s_2^\delta, \dots, s_{2[(n-2)-t_\delta]+3}^\delta \rangle) = \langle v_{p^{j_\delta}(t_\delta)}, v_0, \underline{v_{p^{j_\delta}(t_\delta+1)}, v_0}, \dots, \underline{v_{p^{j_\delta}(n-2)}, v_0}, v_n \rangle \text{ for some } j_\delta \in \{1, \dots, n-2\} \text{ and } t_\delta \in \{1, \dots, n-2\}\}$, $D_2 = \{\delta \in D : \beta(s_1^\delta) = v_n\}$, and $D_3 = \{\delta \in D : s_2^\delta \text{ exists and } \bar{\beta}(\langle s_1^\delta, s_2^\delta, \dots, s_{2t_\delta+1}^\delta \rangle) = \langle v_{p^{j_\delta}(t_\delta)}, v_0, \underline{v_{p^{j_\delta}(t_\delta-1)}, v_0}, \dots, \underline{v_{p^{j_\delta}(1)}, v_0}, v_{n-1} \rangle \text{ for some } j_\delta \in \{1, \dots, n-2\} \text{ and } t_\delta \in \{1, \dots, n-2\}\}$. Then, $D_i \cap D_j = \emptyset$ for each distinct $i, j \in \{0, \dots, 3\}$. Without loss of generality, suppose D_0 is empty or there exists $\tilde{\delta} \in \{1, \dots, \tilde{n}\}$ so that $D_0 = \{\tilde{\delta}, \tilde{\delta} + 1, \dots, \tilde{n}\}$. By Proposition 4.12, $\bigcup_{i=0}^3 D_i = D$, and j_δ and t_δ are unique for each $\delta \in D_1 \cup D_3$. If $\delta \in D_2$, let j_δ be the unique integer j guaranteed by the conclusions of Propositions 4.15 and 4.14.

Let $\tilde{k}_\delta = k_\delta + 2(t_\delta - 1)$ if $\delta \in D_1$, $\tilde{k}_\delta = k_\delta + 2(n - 2)$ if $\delta \in D_2$, $\tilde{k}_\delta = k_\delta - 2t_\delta$ if $\delta \in D_3$, and $\tilde{k}_\delta = k_\delta$ if $\delta \in \{1, \dots, \tilde{n}\} \setminus D$. Let $\tilde{n}_0 = \tilde{n} - |D_0|$ and $\tilde{S}^1, \tilde{S}^2, \dots, \tilde{S}^{\tilde{n}_0}$ be $\tilde{n}_0$ arcs so that $\tilde{S}^i = \langle \tilde{s}_0, \tilde{s}_1^i, \dots, \tilde{s}_{\tilde{k}_i}^i \rangle$ for each $i \in \{1, \dots, \tilde{n}_0\}$ and $\tilde{S}^i \cap \tilde{S}^j = \{\tilde{s}_0\}$ for each distinct $i, j \in \{1, \dots, \tilde{n}_0\}$. Let $\tilde{T}$ be a simple- $\tilde{n}_0$ -od so that $\bigcup_{i=1}^{\tilde{n}_0} \tilde{S}^i = \tilde{T}$ and $\tilde{\beta} : \tilde{T} \rightarrow {}_nX_0$ be the simplicial map defined as: $\tilde{\beta}(\tilde{s}_0) = \beta(s_0)$, $\tilde{\beta}(\langle \tilde{s}_1^\delta, \tilde{s}_2^\delta, \dots, \tilde{s}_{2(t_\delta-1)}^\delta \rangle) = \langle v_{p^{j_\delta}(1)}, v_0, \underline{v_{p^{j_\delta}(2)}, v_0}, \dots, \underline{v_{p^{j_\delta}(t_\delta-1)}, v_0} \rangle$ if $\delta \in D_1$ and $t_\delta > 1$, $\tilde{\beta}(\tilde{s}_k^\delta) = \beta(s_{k-2(t_\delta-1)}^\delta)$ for $k \in \{2(t_\delta - 1) + 1, \dots, \tilde{k}_\delta\}$ if $\delta \in D_1$, $\tilde{\beta}(\langle \tilde{s}_1^\delta, \tilde{s}_2^\delta, \dots, \tilde{s}_{2(n-2)}^\delta \rangle) = \langle v_{p^{j_\delta}(1)}, v_0, \underline{v_{p^{j_\delta}(2)}, v_0}, \dots, \underline{v_{p^{j_\delta}(n-2)}, v_0} \rangle$ if $\delta \in D_2$, $\tilde{\beta}(\tilde{s}_k^\delta) = \beta(s_{k-2(n-2)}^\delta)$ for $k \in \{2(n - 2) + 1, \dots, \tilde{k}_\delta\}$ if $\delta \in D_2$, $\tilde{\beta}(\tilde{s}_k^\delta) = \beta(s_{k+2t_\delta}^\delta)$ for $k \in \{1, \dots, \tilde{k}_\delta\}$ if $\delta \in D_3$, and $\tilde{\beta}(\tilde{s}_k^\delta) = \beta(s_k^\delta)$ for $k \in \{1, \dots, \tilde{k}_\delta\}$ if $\delta \in \{1, \dots, \tilde{n}\} \setminus D$.

Let $E = \{\langle x_0, x_1, \dots, x_{2(n-2)} \rangle \subseteq {}_nX_m : x_0, x_1, \dots, x_{2(n-2)} \in V(nX_m) \text{ are distinct and } \overline{{}_n\Phi_0^m}(\langle x_0, x_1, \dots, x_{2(n-2)+1} \rangle) = \langle v_0, v_{p^{j_e}(1)}, v_0, \underline{v_{p^{j_e}(2)}, v_0}, \dots, \underline{v_{p^{j_e}(n-2)}, v_0}, v_n \rangle \text{ for some } j_e \in \{1, \dots, n-2\} \text{ and some } x_{2(n-2)+1} \in V(nX_m) \text{ adjacent to } x_{2(n-2)}\}$. By Proposition 4.12, j_e is unique for each $e \in E$, and if $e_1, e_2 \in E$ are distinct, ${}_nX_m \supseteq e_1 \cap e_2 = \emptyset$.

Let $e \in E$. By Proposition 4.12, there exist $i \in \{1, \dots, n\}$ and $r \in \{0, \dots, \ell_i\}$ so that

$$\text{a) } \langle x_0, x_1, \dots, x_{2(n-2)} \rangle = \langle x_r^i, x_{r+1}^i, \dots, x_{r+2(n-2)}^i \rangle \text{ or}$$

$$\text{b) } \langle x_0, x_1, \dots, x_{2(n-2)} \rangle = \langle x_r^i, x_{r-1}^i, \dots, x_{r-2(n-2)}^i \rangle.$$

If (a), let $\delta_e \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{r+2(n-2)+1}^i) \in S^{\delta_e} \setminus \{s_0\}$ and $t_e \in \{0, \dots, 2(n-2) + 1\}$ so that $t_e = \min\{t \in \{0, \dots, 2(n-2) + 1\} : \alpha(x_{r+t}^i) \in S^{\delta_e} \setminus \{s_0\}\}$. If (b), let $\delta_e \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_{r-2(n-2)-1}^i) \in S^{\delta_e} \setminus \{s_0\}$ and $t_e \in \{0, \dots, 2(n-2) + 1\}$ so that $t_e = \min\{t \in \{0, \dots, 2(n-2) + 1\} : \alpha(x_{r-t}^i) \in S^{\delta_e} \setminus \{s_0\}\}$. Let $E_0 = \{e \in E : t_e > 1\}$.

Let $\tilde{\alpha} : {}_nX_m \longrightarrow \tilde{T}$ be defined as: $\tilde{\alpha}(x) = \tilde{s}_0$ if $x = x_0$ where $x_0 \in e$ for some $e \in E_0$, $\tilde{\alpha}(x) = \tilde{s}_t^{\delta_e}$ if $x = x_t$ where $t > 0$ and $x_t \in e$ for some $e \in E_0$, $\tilde{\alpha}(x) = \tilde{s}_0$ if $x \in V({}_nX_m \setminus \bigcup E_0)$ with $\alpha(x) = s_0$, $\tilde{\alpha}(x) = \tilde{s}_{k+2(t_\delta-1)}^\delta$ if $x \in V({}_nX_m \setminus \bigcup E_0)$ with $\alpha(x) = s_k^\delta$ and $\delta \in D_1$, $\tilde{\alpha}(x) = \tilde{s}_{k+2(n-2)}^\delta$ if $x \in V({}_nX_m \setminus \bigcup E_0)$ with $\alpha(x) = s_k^\delta$ and $\delta \in D_2$, $\tilde{\alpha}(x) = \tilde{s}_0$ if $x \in V({}_nX_m \setminus \bigcup E_0)$ with $\alpha(x) = s_{2t_\delta}^\delta$ and $\delta \in D_3$, $\tilde{\alpha}(x) = \tilde{s}_{k-2t_\delta}^\delta$ if $x \in V({}_nX_m \setminus \bigcup E_0)$ with $\alpha(x) = s_k^\delta$ where $k > 2t_\delta$ and $\delta \in D_3$, and $\tilde{\alpha}(x) = \tilde{s}_k^\delta$ if $x \in V({}_nX_m \setminus \bigcup E_0)$ with $\alpha(x) = s_k^\delta$ and $\delta \in \{1, \dots, \tilde{n}\} \setminus D$.

CLAIM: $\tilde{\alpha}$ is well-defined.

Let $e \in E_0$. Then, $\delta_e \in D_1 \cup D_2$, $\tilde{k}_{\delta_e} = k_{\delta_e} + 2(t_{\delta_e} - 1) \geq 2[(n-2) - t_{\delta_e}] + 3 + 2(t_{\delta_e} - 1) = 2[(n-2) - (t_{\delta_e} - 1)] + 1 + 2(t_{\delta_e} - 1) = 2(n-2) + 1$ if $\delta_e \in D_1$, $\tilde{k}_{\delta_e} = k_{\delta_e} + 2(n-2) \geq 1 + 2(n-2)$ if $\delta_e \in D_2$, and so, $\tilde{s}_t^{\delta_e}$ exists for all $t \in \{1, \dots, 2(n-2)\}$.

Let $x \in V({}_nX_m \setminus \bigcup E_0)$, $\delta \in D_1$, and $k \in \{1, \dots, k_\delta\}$ so that $\alpha(x) = s_k^\delta$. Then, $k + 2(t_\delta - 1) \leq k_\delta + 2(t_\delta - 1) = \tilde{k}_\delta$, and so, $s_{k+2(t_\delta-1)}^\delta$ exists.

Let $x \in V({}_nX_m \setminus \bigcup E_0)$, $\delta \in D_2$, and $k \in \{1, \dots, k_\delta\}$ so that $\alpha(x) = s_k^\delta$. Then, $k + 2(n-2) \leq k_\delta + 2(n-2) = \tilde{k}_\delta$, and so, $s_{k+2(n-2)}^\delta$ exists.

Let $x \in V({}_nX_m)$, $\delta \in D_3$, and $k \in \{1, \dots, 2t_\delta - 1\}$ so that $\alpha(x) = s_k^\delta$. By Proposition 4.12, $x \in e$ for some $e \in E_0$, and so, $\tilde{\alpha}(x)$ is defined. Thus, if $x \in V({}_nX_m \setminus \bigcup E_0)$, $\delta \in D_3$, and $k \in \{1, \dots, k_\delta\}$ so that $\alpha(x) = s_k^\delta$, then $k \in \{2t_\delta, \dots, k_\delta\}$.

Let $x \in V({}_nX_m \setminus \bigcup E_0)$, $\delta \in \{1, \dots, \tilde{n}\} \setminus D$, and $k \in \{1, \dots, k_\delta\}$ so that $\alpha(x) = s_k^\delta$. Then $k \leq k_\delta = \tilde{k}_\delta$, and so, $\tilde{s}_k^\delta$ exists.

Let $x \in V({}_nX_m)$ and $\delta \in D_0$ so that $\alpha(x) = s_1^\delta$. Then, ${}_n\Phi_0^m(x) \in \{v_i : i \in \{1, \dots, n-2\}\}$. By Proposition 4.12 and definition of D_0 , $x \in e$ for some $e \in E_0$, and so, $\tilde{\alpha}(x)$ is defined.

CLAIM: $\tilde{\alpha}$ is simplicial.

Let $x, y \in V({}_nX_m)$ be distinct and adjacent.

Case 1: $x, y \in e$ for some $e \in E_0$.

Then, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent by definition of $\tilde{\alpha}$ on e .

Case 2: $x \in e$ for some $e \in E_0$ and $y \in V({}_nX_m \setminus \bigcup E_0)$.

Then, $x = x_0$ or $x = x_{2(n-2)}$ where $e = \langle x_0, x_1, \dots, x_{2(n-2)} \rangle$.

Case 2.a: $x = x_0$.

By Proposition 4.12, ${}_n\Phi_0^m(y) = v_{n-1}$. Let $\delta'_e \in \{1, \dots, \tilde{n}\}$ so that $\alpha(x_1) \in S^{\delta'_e} \setminus \{s_0\}$.

Case 2.a.i: $\alpha(x_0) = s_0$.

Then, $\alpha(y) = s_1^\delta$ for some $\delta \in \{1, \dots, \tilde{n}\}$, and since $\beta(s_1^\delta) = \beta(\alpha(y)) = {}_n\Phi_0^m(y) = v_{n-1}$, $\delta \notin D$. So, $\tilde{\alpha}(y) = \tilde{s}_1^\delta$, and since $\tilde{\alpha}(x_0) = \tilde{s}_0$, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent.

Case 2.a.ii: $\alpha(x_0) = s_{k+1}^{\delta'_e}$, where $\alpha(x_1) = s_k^{\delta'_e}$ for some $k \in \{1, \dots, k_{\delta'_e}\}$.

Then, $\alpha(y) = s_{k+2}^{\delta'_e}$ and $\delta'_e \in D_3$, giving $k+2 = 2t_{\delta'_e} + 1$, and so, $\tilde{\alpha}(y) = s_1^{\delta'_e}$. Since $\tilde{\alpha}(x_0) = \tilde{s}_0$, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent.

Since ${}_n\Phi_0^m(y) = v_{n-1}$, by definition of e and E_0 , these are all of the subcases of case 2.a.

Case 2.b: $x = x_{2(n-2)}$.

By definition of E , $y = x_{2(n-2)+1}$ and ${}_n\Phi_0^m(y) = v_n$. Let $\delta_e \in \{1, \dots, \tilde{n}\}$ and $k \in \{1, \dots, k_{\delta_e}\}$ so that $\alpha(x_{2(n-2)+1}) = s_k^{\delta_e}$. Then, by definition of e and E_0 , $\delta_e \in D_1 \cup D_2$.

Case 2.b.i: $\delta_e \in D_1$.

Then, $k = 2[(n-2) - t_{\delta_e}] + 3$, giving $\tilde{\alpha}(y) = \tilde{s}_{k+2(t_{\delta_e}-1)}^{\delta_e} = \tilde{s}_{2(n-2)+1}^{\delta_e}$. Since $\tilde{\alpha}(x_{2(n-2)}) = \tilde{s}_{2(n-2)}^{\delta_e}$, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent.

Case 2.b.ii: $\delta_e \in D_2$.

Then, $k = 1$, giving $\tilde{\alpha}(y) = \tilde{s}_{1+2(n-2)}^{\delta_e}$, and since $\tilde{\alpha}(x_{2(n-2)}) = \tilde{s}_{2(n-2)}^{\delta_e}$, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent.

Case 3: $x, y \in V({}_nX_m \setminus \bigcup E_0)$.

Case 3.a: $\alpha(x), \alpha(y) \in S^\delta \setminus \{s_0\}$ for some $\delta \in \{1, \dots, \tilde{n}\} \setminus D_0$.

Since α is simplicial, then $\alpha(x)$ and $\alpha(y)$ are adjacent by Proposition 4.12, and so, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent by definition of $\tilde{\alpha}$.

Case 3.b: $\alpha(x) = s_0$.

Case 3.b.i: $\alpha(y) = s_1^\delta$ for some $\delta \in \{1, \dots, \tilde{n}\} \setminus D$.

Then, $\tilde{\alpha}(y) = \tilde{s}_1^\delta$ and $\tilde{\alpha}(x) = \tilde{s}_0$, and so, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent.

Case 3.b.ii: $\alpha(y) = s_1^\delta$ for some $\delta \in D_1$.

Then, ${}_n\Phi_0^m(y) = \beta(\alpha(y)) = \beta(s_1^\delta) = v_{p^{j_\delta}(t_\delta)}$. Since $y \in V({}_nX_m \setminus \bigcup E_0)$ and by Proposition 4.12, $y \in e$ for some $e \in E$, then $t_\delta = 1$. Since $\tilde{\alpha}(y) = \tilde{s}_{1+2(t_\delta-1)}^\delta$, then $\tilde{\alpha}(y) = \tilde{s}_{1+2(1-1)}^\delta = s_1^\delta$, and so, $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent since $\tilde{\alpha}(x) = \tilde{s}_0$.

By Propositions 4.12, 4.15, and 4.14, these are all of the subcases of Case 3.b.

CLAIM: $\tilde{\beta}(\tilde{s}_0) = v_0$.

By definition of $\tilde{\beta}$, $\tilde{\beta}(\tilde{s}_0) = \beta(s_0)$, and by definition of β , $\beta(s_0) = v_0$.

CLAIM: $\tilde{\beta} \circ \tilde{\alpha} = {}_n\Phi_0^m$.

Let $x \in V({}_nX_m)$.

Case 1: $x = x_0$ where $x_0 \in e$ for some $e \in E_0$.

Then, ${}_n\Phi_0^m(x) = v_0 = \beta(s_0) = \tilde{\beta}(\tilde{s}_0) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 2: $x = x_t$ where $t > 0$ and $x_t \in e$ for some $e \in E_0$.

Then, $\delta_e \in D_1 \cup D_2$, $j = j_{\delta_e}$ where ${}_n\Phi_0^m(e) = \langle v_0, v_{p^j(1)}, v_0, \underline{v_{p^j(2)}}, v_0, \dots, \underline{v_{p^j(n-2)}}, \underline{v_0} \rangle$, $t_e = 2t_{\delta_e} - 1$ if $\delta_e \in D_1$, and $t_e = 2(n-2) + 1$ if $\delta_e \in D_2$.

Case 2.a: $t \geq t_e$.

Then, $\delta_e \notin D_2$.

Case 2.a.i: t is even.

Since t_e is odd, $t - t_e + 1$ is even and $1 = t_e - t_e + 1 \leq t - t_e + 1 = t - (2t_{\delta_e} - 1) + 1 = t - 2t_{\delta_e} + 2 \leq 2(n-2) - 2t_{\delta_e} + 2 = 2[(n-2) - (t_{\delta_e} - 1)]$. Then, $\beta(s_{t-t_e+1}^{\delta_e}) = v_0$, giving ${}_n\Phi_0^m(x) = v_0 = \beta(s_{t-t_e+1}^{\delta_e}) = \beta(s_{t-2t_{\delta_e}+2}^{\delta_e}) = \beta(s_{t-2(t_{\delta_e}-1)}^{\delta_e}) = \tilde{\beta}(\tilde{s}_t^{\delta_e}) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 2.a.ii: t is odd.

Since t_e is odd, $t - t_e + 1$ is odd and $1 = t_e - t_e + 1 \leq t - t_e + 1 = t - (2t_{\delta_e} - 1) + 1 = t - 2t_{\delta_e} + 2 \leq 2(n - 2) - 1 - 2t_{\delta_e} + 2 = 2[(n - 2) - (t_{\delta_e} - 1)] - 1$. Then, $\beta(s_{t-t_e+1}^{\delta_e}) = v_{p^{j\delta_e}(t_{\delta_e} + \frac{t-t_e+1-1}{2})}$, giving ${}_n\Phi_0^m(x) = v_{p^j(\frac{t+1}{2})} = v_{p^{j\delta_e}(t_{\delta_e} - t_{\delta_e} + \frac{t+1}{2})} = v_{p^{j\delta_e}(t_{\delta_e} - \frac{(2t_{\delta_e}-1)-t}{2})} = v_{p^{j\delta_e}(t_{\delta_e} + \frac{t-t_e}{2})} = v_{p^{j\delta_e}(t_{\delta_e} + \frac{t-t_e+1-1}{2})} = \beta(s_{t-t_e+1}^{\delta_e}) = \beta(s_{t-2t_{\delta_e}+2}^{\delta_e}) = \beta(s_{t-2(t_{\delta_e}-1)}^{\delta_e}) = \tilde{\beta}(\tilde{s}_t^{\delta_e}) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 2.b: $t < t_e$.

Then, $1 \leq t \leq t_e - 1 = 2t_{\delta_e} - 2 = 2(t_{\delta_e} - 1)$ if $\delta_e \in D_1$ and $1 \leq t \leq t_e - 1 = 2(n - 2) + 1 - 1 = 2(n - 2)$ if $\delta_e \in D_2$.

Case 2.b.i: t is even.

Then, ${}_n\Phi_0^m(x) = v_0 = \tilde{\beta}(\tilde{s}_t^{\delta_e}) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 2.b.ii: t is odd.

Then, ${}_n\Phi_0^m(x) = v_{p^j(\frac{t+1}{2})} = v_{p^{j\delta_e}(\frac{t+1}{2})} = \tilde{\beta}(\tilde{s}_t^{\delta_e}) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 3: $x \in V({}_nX_m \setminus \bigcup E_0)$ and $\alpha(x) = s_0$.

Then, ${}_n\Phi_0^m(x) = \beta(\alpha(x)) = \beta(s_0) = \tilde{\beta}(\tilde{s}_0) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 4: $x \in V({}_nX_m \setminus \bigcup E_0)$ and $\alpha(x) = s_k^\delta$ for some $\delta \in D_1 \cup D_2$.

Then, ${}_n\Phi_0^m(x) = \beta(\alpha(x)) = \beta(s_k^\delta) = \tilde{\beta}(\tilde{s}_{k+2(t_{\delta}-1)}^\delta) = \tilde{\beta}(\tilde{\alpha}(x))$ if $\delta \in D_1$ and ${}_n\Phi_0^m(x) = \beta(\alpha(x)) = \beta(s_k^\delta) = \tilde{\beta}(\tilde{s}_{k+2(n-2)}^\delta) = \tilde{\beta}(\tilde{\alpha}(x))$ if $\delta \in D_2$.

Case 5: $x \in V({}_nX_m \setminus \bigcup E_0)$ and $\alpha(x) = s_k^\delta$ for some $\delta \in D_3$.

Then, ${}_n\Phi_0^m(x) = \beta(\alpha(x)) = \beta(s_k^\delta) = \tilde{\beta}(\tilde{s}_{k-2t_\delta}^\delta) = \tilde{\beta}(\tilde{\alpha}(x))$.

Case 6: $x \in V({}_nX_m \setminus \bigcup E_0)$ and $\alpha(x) = s_k^\delta$ for some $\delta \in \{1, \dots, \tilde{n}\} \setminus D$.

Then, ${}_n\Phi_0^m(x) = \beta(\alpha(x)) = \beta(s_k^\delta) = \tilde{\beta}(\tilde{s}_k^\delta) = \tilde{\beta}(\tilde{\alpha}(x))$.

CLAIM: If $x, y, z \in V({}_nX_m)$ so that x, y, z are distinct, $\langle x, y, z \rangle \subseteq {}_nX_m$, and $\tilde{\alpha}(y) = \tilde{s}_0$, then ${}_n\Phi_0^m(x) = v_{n-1}$ or ${}_n\Phi_0^m(z) = v_{n-1}$.

Suppose ${}_n\Phi_0^m(x) \neq v_{n-1}$. By Proposition 4.12, since ${}_n\Phi_0^m(y) = \tilde{\beta}(\tilde{\alpha}(y)) = \tilde{\beta}(\tilde{s}_0) = \beta(s_0) = v_0$, ${}_n\Phi_0^m(x) \neq v_0$ and ${}_n\Phi_0^m(z) \neq v_0$, and so, there exists $\delta \in \{1, \dots, \tilde{n}\}$ so that $\tilde{\alpha}(x) = s_1^\delta$. By Proposition 4.12, $\langle x, y, z \rangle \subseteq {}_nA_m^i$ for some $i \in \{1, \dots, \tilde{n}\}$.

Case 1: $\delta \in D_1 \cup D_2$.

Case 1.a: $x \in e$ for some $e \in E_0$.

Then, $\delta_e = \delta$, $x = x_1$, and $y = x_0$ where $x_0, x_1 \in e$, and so by Proposition 4.12, ${}_n\Phi_0^m(z) = v_{n-1}$.

Case 1.b: $x \in V({}_nX_m \setminus \bigcup E_0)$.

Then $\delta \notin D_2$. Since $\tilde{\alpha}(x) = \tilde{s}_{k+2(t_\delta-1)}^\delta$ where $\alpha(x) = s_k^\delta$, $\tilde{\alpha}(x) = \tilde{s}_1^\delta$, and $1 \leq k \leq k + 2(t_\delta - 1) = 1$, then $k = 1$ and $t_\delta = 1$, giving $\alpha(x) = s_1^\delta$. If $\alpha(y) = s_2^\delta$ and $y \in e$ for some $e \in E_0$, then $x \in e$, giving a contradiction. Thus, $\alpha(y) = s_0$. By Proposition 4.12, since ${}_n\Phi_0^m(x) \neq v_{n-1}$, there exists $\delta' \in \{1, \dots, \tilde{n}\} \setminus \{\delta\}$ so that $\alpha(z) = s_1^{\delta'}$. If ${}_n\Phi_0^m(z) \neq v_{n-1}$, then $x \in e$ for some $e \in E_0$ by Proposition 4.12, resulting in a contradiction. Thus, ${}_n\Phi_0^m(z) = v_{n-1}$.

Case 2: $\delta \in D_3$.

Then, ${}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_1^\delta) = \beta(s_{1+2t_\delta}^\delta) = v_{n-1}$, giving a contradiction. Thus, case 2 does not occur.

Case 3: $\delta \in \{1, \dots, \tilde{n}\} \setminus D$.

Case 3.a: $x \in e$ for some $e \in E_0$.

Then, $\delta_e = \delta$ and $\delta_e \in D_1 \cup D_2$, giving a contradiction. Thus, Case 3.a does not occur.

Case 3.b: $x \in V({}_nX_m) \setminus \bigcup E_0$.

Then, $\alpha(x) = s_k^\delta$, where $\tilde{\alpha}(x) = \tilde{s}_k^\delta$, and so, $\alpha(x) = s_1^\delta$. If $\alpha(y) = s_2^\delta$ and $y \in e$ for some $e \in E_0$, then $\delta \in D$, resulting in a contradiction. Thus, $\alpha(y) = s_0$. Since ${}_n\Phi_0^m(x) \neq v_{n-1}$ and $\delta \notin D$, then ${}_n\Phi_0^m(z) = v_{n-1}$. $\square$

Proposition 4.17. *If $m, \tilde{n}, T, \alpha$, and β exist, then for some $\tilde{n}_1 \in \{1, \dots, \tilde{n}_0\}$ where $\tilde{n}_0$ is as given in Proposition 4.16, there exist a simple- $\tilde{n}_1$ -od $\hat{T}$, $\tilde{n}_1$ arcs $\hat{S}^1, \hat{S}^2, \dots, \hat{S}^{\tilde{n}_1}$ so that $\bigcup_{i=1}^{\tilde{n}_1} \hat{S}^i = \hat{T}$, $\hat{S}^i \cap \hat{S}^j = \{\hat{s}_0\}$ for each distinct $i, j \in \{1, \dots, \tilde{n}_1\}$, so $\hat{s}_0$ is an endpoint of $\hat{S}^i$ for each $i \in \{1, \dots, \tilde{n}_1\}$, and $\hat{S}^i = \langle \hat{s}_0, \hat{s}_1^i, \dots, \hat{s}_{\hat{k}_i}^i \rangle$ for some $\hat{k}_i \in \{1, 2, \dots\}$ for each $i \in \{1, \dots, \tilde{n}_1\}$, and simplicial maps $\hat{\alpha} : {}_nX_m \longrightarrow \hat{T}$ and $\hat{\beta} : \hat{T} \longrightarrow {}_nX_0$ so that $\hat{\beta} \circ \hat{\alpha} = {}_n\Phi_0^m$, $\hat{\beta}(\hat{s}_0) = v_0$, and $\hat{\beta}(\hat{s}_1^\delta) = v_{n-1}$ for each $\delta \in \{1, \dots, \tilde{n}_1\}$.*

Proof. Let $\tilde{T}, \tilde{\alpha}$, and $\tilde{\beta}$ be as given in Proposition 4.16.

Let $F_0 = \{\delta \in \{1, \dots, \tilde{n}_0\} : \tilde{\beta}(\tilde{s}_1^\delta) \neq v_{n-1}\}$, $F_1 = \{\delta \in \{1, \dots, \tilde{n}_0\} : \tilde{s}_3^\delta \text{ exists and } \overline{\tilde{\beta}}(\langle \tilde{s}_1^\delta, \tilde{s}_2^\delta, \tilde{s}_3^\delta \rangle) = \langle v_{n-1}, v_{n+1}, v_{n-1} \rangle, \text{ or } \tilde{s}_3^\delta \text{ exists and } \overline{\tilde{\beta}}(\langle \tilde{s}_1^\delta, \tilde{s}_2^\delta, \tilde{s}_3^\delta \rangle) = \langle v_{n-1}, v_0, v \rangle \text{ for some } v \in \{v_1, v_2, \dots, v_n\} \setminus \{v_{n-1}\}\}$, $F_2 = \{\delta \in \{1, \dots, \tilde{n}_0\} : \tilde{s}_3^\delta \text{ exists and } \overline{\tilde{\beta}}(\langle \tilde{s}_1^\delta, \tilde{s}_2^\delta, \tilde{s}_3^\delta \rangle) = \langle v_{n-1}, v_0, v_{n-1} \rangle\}$, $F_3 = \{\delta \in \{1, \dots, \tilde{n}_0\} : \tilde{s}_3^\delta \text{ does not exist and } \tilde{\beta}(\tilde{s}_1^\delta) = v_{n-1}\}$, and $F = \bigcup_{i=0}^3 F_i$. Then, $F_i \cap F_j = \emptyset$ for each distinct $i, j \in \{0, \dots, 3\}$, and by Proposition 4.12, $|F| = \tilde{n}_0$. Without loss of generality, suppose F_3 is empty or there exists $\tilde{\delta} \in \{1, \dots, \tilde{n}_0\}$ so that $F_3 = \{\tilde{\delta}, \tilde{\delta} + 1, \dots, \tilde{n}_0\}$.

Let $\hat{k}_\delta = \tilde{k}_\delta + 2$ if $\delta \in F_0$, $\hat{k}_\delta = \tilde{k}_\delta$ if $\delta \in F_1$, and $\hat{k}_\delta = \tilde{k}_\delta - 2$ if $\delta \in F_2$. Let $\tilde{n}_1 = \tilde{n}_0 - |F_3|$ and $\hat{S}^1, \hat{S}^2, \dots, \hat{S}^{\tilde{n}_1}$ be $\tilde{n}_1$ arcs so that $\hat{S}^i = \langle \hat{s}_0, \hat{s}_1^i, \dots, \hat{s}_{\hat{k}_i}^i \rangle$ for each $i \in \{1, \dots, \tilde{n}_1\}$ and $\hat{S}^i \cap \hat{S}^j = \{\hat{s}_0\}$ for each distinct $i, j \in \{1, \dots, \tilde{n}_1\}$. Let $\hat{T}$ be a simple- $\tilde{n}_1$ -od so that $\bigcup_{i=1}^{\tilde{n}_1} \hat{S}^i = \hat{T}$ and $\hat{\beta} : \hat{T} \rightarrow {}_nX_0$ be the simplicial map defined as: $\hat{\beta}(\hat{s}_0) = \tilde{\beta}(\tilde{s}_0)$, $\overline{\hat{\beta}}(\langle \hat{s}_1^\delta, \hat{s}_2^\delta \rangle) = \langle v_{n-1}, v_0 \rangle$ if $\delta \in F_0$, $\hat{\beta}(\hat{s}_k^\delta) = \tilde{\beta}(\tilde{s}_{k-2}^\delta)$ for $k \in \{3, \dots, \hat{k}_\delta\}$ if $\delta \in F_0$, $\hat{\beta}(\hat{s}_k^\delta) = \tilde{\beta}(\tilde{s}_k^\delta)$ for $k \in \{1, \dots, \hat{k}_\delta\}$ if $\delta \in F_1$, and $\hat{\beta}(\hat{s}_k^\delta) = \tilde{\beta}(\tilde{s}_{k+2}^\delta)$ for $k \in \{1, \dots, \hat{k}_\delta\}$ if $\delta \in F_2$.

Let $G_0 = \{\langle x_0^i, x_1^i, x_2^i \rangle : i \in \{1, \dots, n\}\}$, $G_1 = \{\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle : \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1}, v_0 \rangle \text{ for some } i \in \{1, \dots, n\} \text{ and for some } r \in \{0, \dots, \ell_i\}\}$, $G_2 = \{\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle : \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1}, v_{n+1} \rangle \text{ for some } i \in \{1, \dots, n\} \text{ and for some } r \in \{0, \dots, \ell_i\}\}$, $G_3 = \{\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle : \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_{n+1}, v_{n-1}, v_0, v_{n-1}, v_0 \rangle \text{ for some } i \in \{1, \dots, n\} \text{ and for some } r \in \{0, \dots, \ell_i\}\}$, $G_4 = \{\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle : \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_{n+1}, v_{n-1}, v_0, v_{n-1}, v_{n+1} \rangle \text{ for some } i \in \{1, \dots, n\} \text{ and for some } r \in \{0, \dots, \ell_i\}\}$, and $G = \bigcup_{i=0}^4 G_i$. Then, $G_i \cap G_j = \emptyset$ for each distinct $i, j \in \{0, \dots, 4\}$.

Let $g_1, g_2 \in G$ be distinct. By Proposition 4.12, if ${}_nX_m \supseteq g_1 \cap g_2 \neq \emptyset$, then $g_1, g_2 \in \bigcup_{i=1}^4 G_i$ with $x_{r_1+4}^{i_1} = x_{r_2}^{i_2}$ and ${}_n\Phi_0^m(x_{r_1+4}^{i_1}) = {}_n\Phi_0^m(x_{r_2}^{i_2}) = v_{n+1}$ (or $x_{r_2+4}^{i_2} = x_{r_1}^{i_1}$ and ${}_n\Phi_0^m(x_{r_1}^{i_1}) = {}_n\Phi_0^m(x_{r_2+4}^{i_2}) = v_{n+1}$), where $g_1 = \langle x_{r_1}^{i_1}, x_{r_1+1}^{i_1}, \dots, x_{r_1+4}^{i_1} \rangle$ and $g_2 = \langle x_{r_2}^{i_2}, x_{r_2+1}^{i_2}, \dots, x_{r_2+4}^{i_2} \rangle$, or $g_1, g_2 \in G_0$ with $x_0^{i_1} = x_0^{i_2} = v_0$, where $g_1 = \langle x_0^{i_1}, x_1^{i_1}, x_2^{i_1} \rangle$ and $g_2 = \langle x_0^{i_2}, x_1^{i_2}, x_2^{i_2} \rangle$.

Let $G_0^H = \{\langle x_0^i, x_1^i, x_2^i \rangle \in G_0 : \tilde{\alpha}(x_k^i) = \tilde{s}_0 \text{ for each } k \in H \text{ and } \tilde{\alpha}(x_k^i) \neq \tilde{s}_0 \text{ for each } k \in \{0, 1, 2\} \setminus H\}$ for $H \subseteq \{0, 1, 2\}$ and $G_j^H = \{\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle \in G_j : \tilde{\alpha}(x_{r+k}^i) = \tilde{s}_0 \text{ for each } k \in H \text{ and } \tilde{\alpha}(x_{r+k}^i) \neq \tilde{s}_0 \text{ for each } k \in \{0, \dots, 4\} \setminus H\}$ for $H \subseteq \{0, \dots, 4\}$ and for $j \in \{1, \dots, 4\}$. Then, if $G_{j_1}^{H_1} \cap G_{j_2}^{H_2} \neq \emptyset$, $H_1 = H_2$ and $j_1 = j_2$.

Let $G'_0 = G_0$ if $G_0^{\{0\}} \neq \emptyset$ or $G_0^{\{0,2\}} \neq \emptyset$, and $G'_0 = \emptyset$ otherwise. Let $G''_0 = G_0$ if $G_0^{\{2\}} \neq \emptyset$, and $G''_0 = \emptyset$ otherwise. Then $\mathcal{G}_1 \cap \mathcal{G}_2 = \emptyset$ for each distinct $\mathcal{G}_1, \mathcal{G}_2 \in \{G_0 \setminus (G' \cup G''), G', G''\}$. Let $G' = G'_0 \cup G''_0 \cup G_1^{\{0\}} \cup G_1^{\{0,2\}} \cup G_1^{\{4\}} \cup G_1^{\{2,4\}} \cup G_1^{\{0,4\}} \cup G_1^{\{0,2,4\}} \cup G_2^{\{0\}} \cup G_2^{\{0,2\}} \cup G_3^{\{4\}} \cup G_3^{\{2,4\}}$.

Let $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+t}^i \rangle \in G$ and $\tilde{t} \in \{0, \dots, t\}$. Let ${}_g\delta_{\tilde{t}} \in \{1, \dots, \tilde{n}_0\}$ so that $\tilde{\alpha}(x_{r+\tilde{t}}^i) \in \tilde{S}^{g\delta_{\tilde{t}}} \setminus \{\tilde{s}_0\}$ whenever $\tilde{\alpha}(x_{r+\tilde{t}}^i) \neq \tilde{s}_0$, ${}_g\delta_{t+1} \in \{1, \dots, \tilde{n}_0\}$ so that $\tilde{\alpha}(x_{r+t+1}^i) \in \tilde{S}^{g\delta_{t+1}} \setminus \{\tilde{s}_0\}$ whenever x_{r+t+1}^i exists and $\tilde{\alpha}(x_{r+t+1}^i) \neq \tilde{s}_0$, and ${}_g\delta_{-1} \in \{1, \dots, \tilde{n}_0\}$ so that $\tilde{\alpha}(x_{r-1}^i) \in \tilde{S}^{g\delta_{-1}} \setminus \{\tilde{s}_0\}$ whenever x_{r-1}^i exists and $\tilde{\alpha}(x_{r-1}^i) \neq \tilde{s}_0$. By Proposition 4.12, x_{r+t+1}^i exists, and if $g \in \bigcup_{i=1}^4 G_i$, x_{r-1}^i exists.

Let $\hat{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle) = \langle \hat{s}_0, \hat{s}_1^{g\delta_3}, \hat{s}_2^{g\delta_3} \rangle$ if $g = \langle x_0^i, x_1^i, x_2^i \rangle \in G'_0 \cup G''_0$, $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle \hat{s}_2^{g\delta_{-1}}, \hat{s}_1^{g\delta_{-1}}, \hat{s}_0, \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle$ if $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_1^{\{0\}} \cup G_1^{\{0,2\}}$, $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle$ if $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_1^{\{4\}} \cup G_1^{\{2,4\}}$, $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle \hat{s}_2^{g\delta_{-1}}, \hat{s}_1^{g\delta_{-1}}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle$ if $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_1^{\{0,4\}} \cup G_1^{\{0,2,4\}}$, $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle \hat{s}_2^{g\delta_{-1}}, \hat{s}_1^{g\delta_{-1}}, \hat{s}_0, \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle$ if $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_2^{\{0\}} \cup G_2^{\{0,2\}}$, $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, x_{r+2}^i, x_{r+3}^i, x_{r+4}^i \rangle) = \langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle$ if $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_3^{\{4\}} \cup G_3^{\{2,4\}}$, $\hat{\alpha}(x) = \hat{s}_0$ if $x \in V(nX_m \setminus \bigcup G')$ and $\tilde{\alpha}(x) = \tilde{s}_0$, where $G' = G_0^{\{0\}} \cup G_0^{\{2\}} \cup G_0^{\{0,2\}} \cup G_1^{\{0\}} \cup G_1^{\{0,2\}} \cup G_1^{\{4\}} \cup G_1^{\{2,4\}} \cup G_1^{\{0,4\}} \cup G_1^{\{0,2,4\}} \cup G_2^{\{0\}} \cup G_2^{\{0,2\}} \cup G_3^{\{4\}} \cup G_3^{\{2,4\}}$, $\hat{\alpha}(x) = \hat{s}_{k+2}^\delta$ if $x \in V(nX_m \setminus \bigcup G')$, $\tilde{\alpha}(x) = \tilde{s}_k^\delta$, and $\delta \in F_0$, $\hat{\alpha}(x) = \hat{s}_k^\delta$ if $x \in V(nX_m \setminus \bigcup G')$, $\tilde{\alpha}(x) = \tilde{s}_k^\delta$, and $\delta \in F_1$, $\hat{\alpha}(x) = \hat{s}_0$ if $x \in V(nX_m \setminus \bigcup G')$, $\tilde{\alpha}(x) = \tilde{s}_2^\delta$, and $\delta \in F_2$, and $\hat{\alpha}(x) = \hat{s}_{k-2}^\delta$ if $x \in V(nX_m \setminus \bigcup G')$, $\tilde{\alpha}(x) = \tilde{s}_k^\delta$, $\delta \in F_2$, and $k \in \{3, \dots, \tilde{k}_\delta\}$.

CLAIM: $\hat{\alpha}$ is well-defined.

Case 1: Let $g = \langle x_0^i, x_1^i, x_2^i \rangle \in G'_0 \cup G''_0$.

By Proposition 4.12, ${}_n\Phi_0^m(x_3^i) = v_n$.

Case1.a: Suppose $g \in G'_0$.

Then,

- i) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_0, \tilde{s}_1^{\delta_3}, \tilde{s}_2^{\delta_3}, \tilde{s}_3^{\delta_3} \rangle$ for some $\delta_3 \in \{1, \dots, \tilde{n}_0\}$ or
- ii) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_0, \tilde{s}_1^{\delta_1}, \tilde{s}_0, \tilde{s}_1^{\delta_3} \rangle$ for some $\delta_1, \delta_3 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_1 \neq \delta_3$.

If (i), then $\delta_3 \in F_1$ and $g\delta_3 = \delta_3$, giving $3 \leq \tilde{k}_{\delta_3} = \hat{k}_{g\delta_3}$, and so, $\hat{s}_1^{g\delta_3}$ and $\hat{s}_2^{g\delta_3}$ exist.

If (ii), then $\delta_3 \in F_0$ and $g\delta_3 = \delta_3$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_3} + 2 = \hat{k}_{g\delta_3}$, and so, $\hat{s}_1^{g\delta_3}$ and $\hat{s}_2^{g\delta_3}$ exist.

Case1.b: Suppose $g \in G''$.

Then,

- i) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_0, \tilde{s}_1^{\delta_3} \rangle$ for some $\delta_0, \delta_3 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_0 \neq \delta_3$,
- ii) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_2^{\delta_3}, \tilde{s}_1^{\delta_3}, \tilde{s}_2^{\delta_3}, \tilde{s}_3^{\delta_3} \rangle$ for some $\delta_3 \in \{1, \dots, \tilde{n}_0\}$, or
- iii) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_2^{\delta_3}, \tilde{s}_3^{\delta_3}, \tilde{s}_4^{\delta_3}, \tilde{s}_5^{\delta_3} \rangle$ for some $\delta_3 \in \{1, \dots, \tilde{n}_0\}$.

If (i), then $\delta_3 \in F_0$ and $g\delta_3 = \delta_3$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_3} + 2 = \hat{k}_{g\delta_3}$, and so, $\hat{s}_1^{g\delta_3}$ and $\hat{s}_2^{g\delta_3}$ exist. If (ii), then $\delta_3 \in F_1$ and $g\delta_3 = \delta_3$, giving $3 \leq \tilde{k}_{\delta_3} = \hat{k}_{g\delta_3}$, and so, $\hat{s}_1^{g\delta_3}$ and $\hat{s}_2^{g\delta_3}$ exist. If (iii), then $\delta_3 \in F_2$ and $g\delta_3 = \delta_3$, giving $3 = 5 - 2 \leq \tilde{k}_{\delta_3} - 2 = \hat{k}_{g\delta_3}$, and so, $\hat{s}_1^{g\delta_3}$ and $\hat{s}_2^{g\delta_3}$ exist.

Case 2: Let $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_1^{\{0\}} \cup G_1^{\{0,2\}}$ (or $G_1^{\{4\}} \cup G_1^{\{2,4\}}$).

By Proposition 4.12, ${}_n\Phi_0^m(x_{r-1}^i), {}_n\Phi_0^m(x_{r+5}^i) \notin \{v_0, v_{n-1}, v_{n+1}, v_{n+2}\}$. Then,

- i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_4}, \tilde{s}_2^{\delta_4}, \tilde{s}_3^{\delta_4}, \tilde{s}_4^{\delta_4}, \tilde{s}_5^{\delta_4} \rangle$ for some $\delta_{-1}, \delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_{-1} \neq \delta_4$ (or $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_5^{\delta_0}, \tilde{s}_4^{\delta_0}, \tilde{s}_3^{\delta_0}, \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_0, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_0 \neq \delta_5$),

- ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_4}, \tilde{s}_2^{\delta_4}, \tilde{s}_1^{\delta_4}, \tilde{s}_2^{\delta_4}, \tilde{s}_3^{\delta_4} \rangle$ for some $\delta_{-1}, \delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_{-1} \neq \delta_4$ (or $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_3^{\delta_0}, \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_0, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_0 \neq \delta_5$), or

- iii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_1}, \tilde{s}_0, \tilde{s}_1^{\delta_4}, \tilde{s}_2^{\delta_4}, \tilde{s}_3^{\delta_4} \rangle$ for some $\delta_{-1}, \delta_1, \delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_{-1} \neq \delta_1$ and $\delta_{-1} \neq \delta_4$ (or $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_3^{\delta_0}, \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_0, \tilde{s}_1^{\delta_3}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_0, \delta_3, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_5 \neq \delta_3$ and $\delta_5 \neq \delta_0$).

If (i), then $\delta_{-1} \in F_0$, $\delta_4 \in F_2$, $g\delta_{-1} = \delta_{-1}$, and $g\delta_4 = \delta_4$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_{-1}} + 2 = \hat{k}_{g\delta_{-1}}$ and $3 = 5 - 2 \leq \tilde{k}_{\delta_4} - 2 = \hat{k}_{g\delta_4}$, and so, $\hat{s}_1^{g\delta_{-1}}$, $\hat{s}_2^{g\delta_{-1}}$, $\hat{s}_1^{g\delta_4}$, and $\hat{s}_2^{g\delta_4}$ exist (or $\delta_0 \in F_2$, $\delta_5 \in F_0$, $g\delta_0 = \delta_0$, and $g\delta_5 = \delta_5$, giving $3 = 5 - 2 \leq \tilde{k}_{\delta_0} - 2 = \hat{k}_{g\delta_0}$

and $3 = 1 + 2 \leq \tilde{k}_{\delta_5} + 2 = \hat{k}_{g\delta_5}$, and so, $\hat{s}_1^{g\delta_0}$, $\hat{s}_2^{g\delta_0}$, $\hat{s}_1^{g\delta_5}$, and $\hat{s}_2^{g\delta_5}$ exist).

If (ii), then $\delta_{-1} \in F_0$, $\delta_4 \in F_1$, $g\delta_{-1} = \delta_{-1}$, and $g\delta_4 = \delta_4$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_{-1}} + 2 = \hat{k}_{g\delta_{-1}}$ and $3 \leq \tilde{k}_{\delta_4} = \hat{k}_{g\delta_4}$, and so, $\hat{s}_1^{g\delta_{-1}}$, $\hat{s}_2^{g\delta_{-1}}$, $\hat{s}_1^{g\delta_4}$, and $\hat{s}_2^{g\delta_4}$ exist (or $\delta_0 \in F_1$, $\delta_5 \in F_0$, $g\delta_0 = \delta_0$, and $g\delta_5 = \delta_5$, giving $3 \leq \tilde{k}_{\delta_0} = \hat{k}_{g\delta_0}$ and $3 = 1 + 2 \leq \tilde{k}_{\delta_5} + 2 = \hat{k}_{g\delta_5}$, and so, $\hat{s}_1^{g\delta_0}$, $\hat{s}_2^{g\delta_0}$, $\hat{s}_1^{g\delta_5}$, and $\hat{s}_2^{g\delta_5}$ exist).

If (iii), then $\delta_{-1} \in F_0$, $\delta_1 \in F_1 \cup F_2$, $\delta_4 \in F_1$, $g\delta_{-1} = \delta_{-1}$, $g\delta_1 = \delta_1$, and $g\delta_4 = \delta_4$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_{-1}} + 2 = \hat{k}_{g\delta_{-1}}$ and $3 \leq \tilde{k}_{\delta_4} = \hat{k}_{g\delta_4}$, and so, $\hat{s}_1^{g\delta_{-1}}$, $\hat{s}_2^{g\delta_{-1}}$, $\hat{s}_1^{g\delta_4}$, and $\hat{s}_2^{g\delta_4}$ exist (or $\delta_0 \in F_1$, $\delta_3 \in F_1 \cup F_2$, $\delta_5 \in F_0$, $g\delta_0 = \delta_0$, $g\delta_3 = \delta_3$, and $g\delta_5 = \delta_5$, giving $3 \leq \tilde{k}_{\delta_0} = \hat{k}_{g\delta_0}$ and $3 = 1 + 2 \leq \tilde{k}_{\delta_5} + 2 = \hat{k}_{g\delta_5}$, and so, $\hat{s}_1^{g\delta_0}$, $\hat{s}_2^{g\delta_0}$, $\hat{s}_1^{g\delta_5}$, and $\hat{s}_2^{g\delta_5}$ exist).

Case 3: Let $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_1^{\{0,4\}} \cup G_1^{\{0,2,4\}}$.

By Proposition 4.12, ${}_n\Phi_0^m(x_{r-1}^i), {}_n\Phi_0^m(x_{r+5}^i) \notin \{v_0, v_{n-1}, v_{n+1}, v_{n+2}\}$, giving

- i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_2}, \tilde{s}_2^{\delta_2}, \tilde{s}_1^{\delta_5}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_{-1}, \delta_2, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_2 \neq \delta_{-1}$ and $\delta_2 \neq \delta_5$, or
- ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_1}, \tilde{s}_0, \tilde{s}_1^{\delta_3}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_{-1}, \delta_1, \delta_3, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_{-1}, \delta_5 \notin \{\delta_1, \delta_3\}$.

Then, $\delta_{-1}, \delta_5 \in F_0$, $g\delta_{-1} = \delta_{-1}$, and $g\delta_5 = \delta_5$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_{-1}} + 2 = \hat{k}_{g\delta_{-1}}$ and $3 = 1 + 2 \leq \tilde{k}_{\delta_5} + 2 = \hat{k}_{g\delta_5}$, and so, $\hat{s}_1^{g\delta_{-1}}$, $\hat{s}_2^{g\delta_{-1}}$, $\hat{s}_1^{g\delta_5}$, and $\hat{s}_2^{g\delta_5}$ exist.

Case 4: Let $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle \in G_2^{\{0\}} \cup G_2^{\{0,2\}}$ (or $G_3^{\{4\}} \cup G_3^{\{2,4\}}$).

By Proposition 4.12, ${}_n\Phi_0^m(x_{r-1}^i) \notin \{v_0, v_{n-1}, v_{n+1}, v_{n+2}\}$ (or ${}_n\Phi_0^m(x_{r+5}^i) \notin \{v_0, v_{n-1}, v_{n+1}, v_{n+2}\}$), giving

- i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+4}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_4}, \tilde{s}_2^{\delta_4}, \tilde{s}_3^{\delta_4}, \tilde{s}_4^{\delta_4} \rangle$ for some $\delta_{-1}, \delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_{-1} \neq \delta_4$ (or $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_4^{\delta_0}, \tilde{s}_3^{\delta_0}, \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_0, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_0 \neq \delta_5$), or
- ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+4}^i \rangle) = \langle \tilde{s}_1^{\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{\delta_1}, \tilde{s}_0, \tilde{s}_1^{\delta_4}, \tilde{s}_2^{\delta_4} \rangle$ for some $\delta_{-1}, \delta_1, \delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_{-1} \neq \delta_1$ and $\delta_{-1} \neq \delta_4$ (or $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_2^{\delta_0}, \tilde{s}_1^{\delta_0}, \tilde{s}_0, \tilde{s}_1^{\delta_3}, \tilde{s}_0, \tilde{s}_1^{\delta_5} \rangle$ for some $\delta_0, \delta_3, \delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $\delta_5 \neq \delta_0$ and $\delta_5 \neq \delta_3$).

If (i), then $\delta_{-1} \in F_0$, $\delta_4 \in F_2$, $g\delta_{-1} = \delta_{-1}$, and $g\delta_4 = \delta_4$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_{-1}} + 2 = \hat{k}_{g\delta_{-1}}$ and $2 = 4 - 2 \leq \tilde{k}_{\delta_4} - 2 = \hat{k}_{g\delta_4}$, and so, $\hat{s}_1^{g\delta_{-1}}$, $\hat{s}_2^{g\delta_{-1}}$, $\hat{s}_1^{g\delta_4}$, and $\hat{s}_2^{g\delta_4}$ exist (or $\delta_0 \in F_2$, $\delta_5 \in F_0$, $g\delta_0 = \delta_0$, and $g\delta_5 = \delta_5$, giving $2 = 4 - 2 \leq \tilde{k}_{\delta_0} - 2 = \hat{k}_{g\delta_0}$

and $3 = 1 + 2 \leq \tilde{k}_{\delta_5} + 2 = \hat{k}_{g\delta_5}$, and so, $\hat{s}_1^{g\delta_0}$, $\hat{s}_2^{g\delta_0}$, $\hat{s}_1^{g\delta_5}$, and $\hat{s}_2^{g\delta_5}$ exist).

If (ii), then $\delta_{-1} \in F_0$, $\delta_4 \in F_1$, $g\delta_{-1} = \delta_{-1}$, and $g\delta_4 = \delta_4$, giving $3 = 1 + 2 \leq \tilde{k}_{\delta_{-1}} + 2 = \hat{k}_{g\delta_{-1}}$ and $2 \leq \tilde{k}_{\delta_4} = \hat{k}_{g\delta_4}$, and so, $\hat{s}_1^{g\delta_{-1}}$, $\hat{s}_2^{g\delta_{-1}}$, $\hat{s}_1^{g\delta_4}$, and $\hat{s}_2^{g\delta_4}$ exist (or $\delta_0 \in F_1$, $\delta_5 \in F_0$, $g\delta_0 = \delta_0$, and $g\delta_5 = \delta_5$, giving $2 \leq \tilde{k}_{\delta_0} = \hat{k}_{g\delta_0}$ and $3 = 1 + 2 \leq \tilde{k}_{\delta_5} + 2 = \hat{k}_{g\delta_5}$, and so, $\hat{s}_1^{g\delta_0}$, $\hat{s}_2^{g\delta_0}$, $\hat{s}_1^{g\delta_5}$, and $\hat{s}_2^{g\delta_5}$ exist).

Case 5: Let $x \in V(nX_m \setminus \bigcup G')$, $\delta \in F_0$, and $k \in \{1, \dots, \tilde{k}_\delta\}$ so that $\tilde{\alpha}(x) = \tilde{s}_k^\delta$.

Then, $k + 2 \leq \tilde{k}_\delta + 2 = \hat{k}_\delta$, and so, $\hat{s}_{k+2}^\delta$ exists.

Case 6: Let $x \in V(nX_m)$ and $\delta \in F_2$ so that $\tilde{\alpha}(x) = \tilde{s}_1^\delta$.

Then, ${}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_1^\delta) = v_{n-1}$, and by Proposition 4.12, $x \in g$ for some $g \in G$.

Suppose $g \notin G_0$, and let $\tilde{t} \in \{0, \dots, 4\}$ so that $x = x_{r+\tilde{t}}^i$, where $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle$. Then, $\tilde{t} \in \{1, 3\}$. Suppose $\tilde{\alpha}(x_r^i) \neq \tilde{s}_0$ and $\tilde{\alpha}(x_{r+4}^i) \neq \tilde{s}_0$.

If $\tilde{t} = 1$, then $\tilde{\alpha}(x_r^i) = \tilde{s}_2^\delta$, giving ${}_n\Phi_0^m(x_r^i) = v_0$, and $\tilde{\alpha}(x_{r-1}^i) = \tilde{s}_1^\delta$ or $\tilde{\alpha}(x_{r-1}^i) = \tilde{s}_3^\delta$. Thus, ${}_n\Phi_0^m(x_{r-1}^i) = v_{n-1}$ and $\overline{{}_n\Phi_0^m}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+3}^i \rangle) = \langle v_{n-1}, v_0, v_{n-1}, v_0, v_{n-1} \rangle$, contradicting Proposition 4.12.

If $\tilde{t} = 3$, then $\tilde{\alpha}(x_{r+4}^i) = \tilde{s}_2^\delta$, giving ${}_n\Phi_0^m(x_{r+4}^i) = v_0$, and $\tilde{\alpha}(x_{r+5}^i) = \tilde{s}_1^\delta$ or $\tilde{\alpha}(x_{r+5}^i) = \tilde{s}_3^\delta$. Thus, ${}_n\Phi_0^m(x_{r+5}^i) = v_{n-1}$ and $\overline{{}_n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+5}^i \rangle) = \langle v_{n-1}, v_0, v_{n-1}, v_0, v_{n-1} \rangle$, contradicting Proposition 4.12.

Thus, $\tilde{\alpha}(x_r^i) = \tilde{s}_0$ or $\tilde{\alpha}(x_{r+4}^i) = \tilde{s}_0$, and so, $g \in G'$, giving that $\hat{\alpha}(x)$ is defined. If $g \in G_0$, then by Proposition 4.12, $x = x_1^i$ and ${}_n\Phi_0^m(x_3^i) = v_n$, where $g = \langle x_0^i, x_1^i, x_2^i \rangle$, and so, $\tilde{\alpha}(x_2^i) \neq \tilde{s}_2^\delta$. Then, $\tilde{\alpha}(x_2^i) = \tilde{s}_0$, giving $g \in G'_0 \cup G''_0 \subseteq G'$, and so, $\hat{\alpha}(x)$ is defined.

Case 6.5: Let $x \in V(nX_m)$ and $\delta \in F_3$ so that $\tilde{\alpha}(x) = \tilde{s}_1^\delta$.

Then, by Proposition 4.12 and definition of F_3 , $x \in g$ for some $g \in G'$, and so, $\hat{\alpha}(x)$ is defined.

Case 7: Let $x \in V(nX_m \setminus \bigcup G')$, $\delta \in F_2$, and $k \in \{1, \dots, \tilde{k}_\delta\}$ so that $\tilde{\alpha}(x) = \tilde{s}_k^\delta$.

From Case 6, $k \in \{2, \dots, \tilde{k}_\delta\}$. If $k = 2$, then $\hat{\alpha}(x)$ is defined. If $k \in \{3, \dots, \tilde{k}_\delta\}$, then $\hat{k}_\delta = \tilde{k}_\delta - 2 \geq k - 2 \geq 1$, and so, $\hat{s}_{k-2}^\delta$ exists.

Case 8: Let $x \in V(nX_m \setminus \bigcup G')$, $\delta \in F_1$, and $k \in \{1, \dots, \tilde{k}_\delta\}$ so that $\tilde{\alpha}(x) = \tilde{s}_k^\delta$.

Then, $1 \leq k \leq \tilde{k}_\delta = \hat{k}_\delta$, and so, $\hat{s}_k^\delta$ exists.

Case 9: Let $g_1 = \langle x_0^{i_1}, x_1^{i_1}, x_2^{i_1} \rangle, g_2 = \langle x_0^{i_2}, x_1^{i_2}, x_2^{i_2} \rangle \in G_0$ so that $i_1 \neq i_2$ and ${}_nX_m \supseteq g_1 \cap g_2 \neq \emptyset$.

Then, $x_0^{i_1} = x_0^{i_2} = v_0$. If $D'_0 \cup D''_0 \neq \emptyset$, then $\hat{\alpha}(x_0^{i_1}) = \hat{\alpha}(x_0^{i_2}) = \hat{s}_0$. If $D'_0 \cup D''_0 = \emptyset$, then $x_0^{i_1} = x_0^{i_2} = x$ for some x as given in case 5, case 7, or case 8.

Case 10: Let $g_1 = \langle x_{r_1}^i, x_{r_1+1}^i, \dots, x_{r_1+4}^i \rangle, g_2 = \langle x_{r_2}^i, x_{r_2+1}^i, \dots, x_{r_2+4}^i \rangle \in \bigcup_{i=1}^4 G_i$ so that $r_1 \neq r_2$ and ${}_nX_m \supseteq g_1 \cap g_2 \neq \emptyset$.

Then, $x_{r_1+4}^i = x_{r_2}^i$ and ${}_n\Phi_0^m(x_{r_1+4}^i) = {}_n\Phi_0^m(x_{r_2}^i) = v_{n+1}$, or $x_{r_1}^i = x_{r_2+4}^i$ and ${}_n\Phi_0^m(x_{r_1}^i) = {}_n\Phi_0^m(x_{r_2+4}^i) = v_{n+1}$. Without loss of generality, suppose $x_{r_1+4}^i = x_{r_2}^i$. Then, $g_1 \in G_2 \cup G_4$ and $g_2 \in G_3 \cup G_4$.

Case 10.a: $g_1, g_2 \in G'$.

Then, $g_1 \in G_2^{\{0\}} \cup G_2^{\{0,2\}}$ and $g_2 \in G_3^{\{4\}} \cup G_3^{\{2,4\}}$, giving

i) $\bar{\alpha}(\langle x_{r_1-1}^i, x_{r_1}^i, \dots, x_{r_1+4}^i \rangle) = \langle \tilde{s}_1^{g_1\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g_1\delta_4}, \tilde{s}_2^{g_1\delta_4}, \tilde{s}_3^{g_1\delta_4}, \tilde{s}_4^{g_1\delta_4} \rangle$ for some $g_1\delta_{-1}, g_1\delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $g_1\delta_{-1} \neq g_1\delta_4$ or

ii) $\bar{\alpha}(\langle x_{r_1-1}^i, x_{r_1}^i, \dots, x_{r_1+4}^i \rangle) = \langle \tilde{s}_1^{g_1\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g_1\delta_1}, \tilde{s}_0, \tilde{s}_1^{g_1\delta_4}, \tilde{s}_2^{g_1\delta_4} \rangle$ for some $g_1\delta_{-1}, g_1\delta_1, g_1\delta_4 \in \{1, \dots, \tilde{n}_0\}$ so that $g_1\delta_{-1} \neq g_1\delta_1$ and $g_1\delta_{-1} \neq g_1\delta_4$, from case 4.

If (i), then $\bar{\alpha}(\langle x_{r_2}^i, x_{r_2+1}^i, \dots, x_{r_2+5}^i \rangle) = \langle \tilde{s}_4^{g_2\delta_0}, \tilde{s}_3^{g_2\delta_0}, \tilde{s}_2^{g_2\delta_0}, \tilde{s}_1^{g_2\delta_0}, \tilde{s}_0, \tilde{s}_1^{g_2\delta_5} \rangle$ for some $g_2\delta_0, g_2\delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $g_2\delta_0 \neq g_2\delta_5$, from case 4. So, $g_1\delta_4 = g_2\delta_0$, giving $\hat{\alpha}(x_{r_1+4}^i) = \hat{s}_2^{g_1\delta_4} = \hat{s}_2^{g_2\delta_0} = \hat{\alpha}(x_{r_2}^i)$.

If (ii), then $\bar{\alpha}(\langle x_{r_2}^i, x_{r_2+1}^i, \dots, x_{r_2+5}^i \rangle) = \langle \tilde{s}_2^{g_2\delta_0}, \tilde{s}_1^{g_2\delta_0}, \tilde{s}_0, \tilde{s}_1^{g_2\delta_3}, \tilde{s}_0, \tilde{s}_1^{g_2\delta_5} \rangle$ for some $g_2\delta_0, g_2\delta_3, g_2\delta_5 \in \{1, \dots, \tilde{n}_0\}$ so that $g_2\delta_5 \neq g_2\delta_0$ and $g_2\delta_5 \neq g_2\delta_3$, from case 4. So, $g_1\delta_4 = g_2\delta_0$, giving $\hat{\alpha}(x_{r_1+4}^i) = \hat{s}_2^{g_1\delta_4} = \hat{s}_2^{g_2\delta_0} = \hat{\alpha}(x_{r_2}^i)$.

Thus, $\hat{\alpha}(x_{r_1+4}^i)$ is unique, since if $g_3 \in G$ so that $g_3 \neq g_1$ and $g_3 \neq g_2$, $x_{r_1+4}^i \notin g_3$.

Case 10.b: $g_1 \in G'$ and $g_2 \notin G'$, or $g_2 \in G'$ and $g_1 \notin G'$.

Since if $g_3 \in G$ so that $g_3 \neq g_1$ and $g_3 \neq g_2$, $x_{r_1+4}^i \notin g_3$, then g_1 is the unique element of G' containing $x_{r_1+4}^i$ or g_2 is the unique element of G' containing $x_{r_2}^i$, and so, $\hat{\alpha}(x_{r_1+4}^i)$ is unique.

Case 10.c: $g_1, g_2 \notin G'$.

Since if $g_3 \in G$ so that $g_3 \neq g_1$ and $g_3 \neq g_2$, $x_{r_1+4}^i \notin g_3$, then, $x_{r_1+4}^i \in V({}_nX_m \setminus$

$\bigcup G'$), and so, $\hat{\alpha}(x_{r_1+4}^i)$ is unique.

CLAIM: $\hat{\alpha}$ is simplicial.

Let $x, y \in V({}_nX_m)$ be distinct and adjacent.

Case 1: $x, y \in g$ for some $g \in G'$.

Then, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent by definition of $\hat{\alpha}$ on g .

Case 2: $x \in g$ for some $g \in G'$ and $y \in V({}_nX_m \setminus \bigcup G')$.

Case 2.a: $g \in G'_0 \bigcup G''_0$.

Then, $x = x_2^i$ and $y = x_3^i$ where $g = \langle x_0^i, x_1^i, x_2^i \rangle$ for some $i \in \{1, \dots, n\}$.

Case 2.a.i: $g \in G'_0$.

From Case 1.a of the previous claim,

i) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_0, \tilde{s}_1^{g\delta_3}, \tilde{s}_2^{g\delta_3}, \tilde{s}_3^{g\delta_3} \rangle$ for some $g\delta_3 \in F_1$ or

ii) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_0, \tilde{s}_1^{g\delta_1}, \tilde{s}_0, \tilde{s}_1^{g\delta_3} \rangle$ for some $g\delta_1 \in F_1 \bigcup F_2$ and for some $g\delta_3 \in F_0$.

If (i), then $\hat{\alpha}(x) = \hat{\alpha}(x_2^i) = \hat{s}_2^{g\delta_3}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_3^i) = \hat{s}_3^{g\delta_3}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (ii), then $\hat{\alpha}(x) = \hat{\alpha}(x_2^i) = \hat{s}_2^{g\delta_3}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_3^i) = \hat{s}_{1+2}^{g\delta_3} = \hat{s}_3^{g\delta_3}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.a.ii: $g \in G''_0$.

From Case 1.b of the previous claim,

i) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_0, \tilde{s}_1^{g\delta_3} \rangle$ for some $g\delta_3 \in F_0$,

ii) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_2^{g\delta_3}, \tilde{s}_1^{g\delta_3}, \tilde{s}_2^{g\delta_3}, \tilde{s}_3^{g\delta_3} \rangle$ for some $g\delta_3 \in F_1$, or

iii) $\bar{\alpha}(\langle x_0^i, x_1^i, x_2^i, x_3^i \rangle) = \langle \tilde{s}_2^{g\delta_3}, \tilde{s}_3^{g\delta_3}, \tilde{s}_4^{g\delta_3}, \tilde{s}_5^{g\delta_3} \rangle$ for some $g\delta_3 \in F_2$.

If (i), then $\hat{\alpha}(x) = \hat{\alpha}(x_2^i) = \hat{s}_2^{g\delta_3}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_3^i) = \hat{s}_{1+2}^{g\delta_3} = \hat{s}_3^{g\delta_3}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (ii), then $\hat{\alpha}(x) = \hat{\alpha}(x_2^i) = \hat{s}_2^{g\delta_3}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_3^i) = \hat{s}_3^{g\delta_3}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (iii), then $\hat{\alpha}(x) = \hat{\alpha}(x_2^i) = \hat{s}_2^{g\delta_3}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_3^i) = \hat{s}_{5-2}^{g\delta_3} = \hat{s}_3^{g\delta_3}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b: $g \in \bigcup_{i=1}^4 G_i$.

Then, $x = x_r^i$ or $x = x_{r+4}^i$ where $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle$.

Case 2.b.i: $g \in G_1^{\{0\}} \cup G_1^{\{0,2\}}$.

From Case 2 of the previous claim,

i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4}, \tilde{s}_3^{g\delta_4}, \tilde{s}_4^{g\delta_4}, \tilde{s}_5^{g\delta_4} \rangle$ for some $g\delta_{-1} \in F_0$

and $g\delta_4 \in F_2$,

ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4}, \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4}, \tilde{s}_3^{g\delta_4} \rangle$ for some $g\delta_{-1} \in F_0$

and $g\delta_4 \in F_1$, or

iii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_1}, \tilde{s}_0, \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4}, \tilde{s}_3^{g\delta_4} \rangle$ for some $g\delta_{-1} \in F_0$, $g\delta_1 \in F_1 \cup F_2$, and $g\delta_4 \in F_1$.

Case 2.b.i.1: $x = x_r^i$.

Then, $y = x_{r-1}^i$.

If (i), (ii), or (iii), then $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_{-1}}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_{1+2}^{g\delta_{-1}} = \hat{s}_3^{g\delta_{-1}}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.i.2: $x = x_{r+4}^i$.

Then, $y = x_{r+5}^i$.

If (i), then $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_4}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_{5-2}^{g\delta_4} = \hat{s}_3^{g\delta_4}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (ii) or (iii), then $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_4}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_3^{g\delta_4}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.ii: $g \in G_1^{\{4\}} \cup G_1^{\{2,4\}}$.

From Case 2 of the previous claim,

i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_5^{g\delta_0}, \tilde{s}_4^{g\delta_0} \tilde{s}_3^{g\delta_0}, \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_0 \in F_2$ and $g\delta_5 \in F_0$,

ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_3^{g\delta_0}, \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_0 \in F_1$ and $g\delta_5 \in F_0$, or

iii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_3^{g\delta_0}, \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_0, \tilde{s}_1^{g\delta_3}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_0 \in F_1$, $g\delta_3 \in F_1 \cup F_2$, and $g\delta_5 \in F_0$.

Case 2.b.ii.1: $x = x_{r+4}^i$.

Then, $y = x_{r+5}^i$.

If (i), (ii), or (iii), then $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_5}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_{1+2}^{g\delta_5} = \hat{s}_3^{g\delta_5}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.ii.2: $x = x_r^i$.

Then, $y = x_{r-1}^i$.

If (i), then $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_0}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_{5-2}^{g\delta_0} = \hat{s}_3^{g\delta_0}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (ii) or (iii), then $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_0}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_3^{g\delta_0}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.iii: $g \in G_1^{\{0,4\}} \cup G_1^{\{0,2,4\}}$.

From Case 3 of the previous claim,

i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_2}, \tilde{s}_2^{g\delta_2}, \tilde{s}_1^{g\delta_2}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_{-1}, g\delta_5 \in F_0$ and $g\delta_2 \in F_1 \cup F_2$ or

ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_1}, \tilde{s}_0, \tilde{s}_1^{g\delta_3}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_{-1}, g\delta_5 \in F_0$ and $g\delta_1, g\delta_3 \in F_1 \cup F_2$.

Case 2.b.iii.1: $x = x_r^i$.

Then, $y = x_{r-1}^i$.

If (i) or (ii), then $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_{-1}}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_{1+2}^{g\delta_{-1}} = \hat{s}_3^{g\delta_{-1}}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.iii.2: $x = x_{r+4}^i$.

Then, $y = x_{r+5}^i$.

If (i) or (ii), then $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_5}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_{1+2}^{g\delta_5} = \hat{s}_3^{g\delta_5}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.iv: $g \in G_2^{\{0\}} \cup G_2^{\{0,2\}}$.

From Case 4 of the previous claim,

i) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+4}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4}, \tilde{s}_3^{g\delta_4}, \tilde{s}_4^{g\delta_4} \rangle$ for some $g\delta_{-1} \in F_0$ and $g\delta_4 \in F_2$ or

ii) $\bar{\alpha}(\langle x_{r-1}^i, x_r^i, \dots, x_{r+4}^i \rangle) = \langle \tilde{s}_1^{g\delta_{-1}}, \tilde{s}_0, \tilde{s}_1^{g\delta_1}, \tilde{s}_0, \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4} \rangle$ for some $g\delta_{-1} \in F_0$, $g\delta_1 \in F_1 \cup F_2$, and $g\delta_4 \in F_1$.

Case 2.b.iv.1: $x = x_r^i$.

Then, $y = x_{r-1}^i$.

If (i) or (ii), then $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_{-1}}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_{1+2}^{g\delta_{-1}} = \hat{s}_3^{g\delta_{-1}}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.iv.2: $x = x_{r+4}^i$.

Then, $y = x_{r+5}^i$.

If (i), then by Proposition 4.12, x_{r+5}^i exists and ${}_n\Phi_0^m(x_{r+5}^i) = v_{n-1}$, and so, $\tilde{\alpha}(x_{r+5}^i) = \tilde{s}_3^{g\delta_4}$ or $\tilde{\alpha}(x_{r+5}^i) = \tilde{s}_5^{g\delta_4}$. Then, $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_4}$, and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_{3-2}^{g\delta_4} = \hat{s}_1^{g\delta_4}$ or $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_{5-2}^{g\delta_4} = \hat{s}_3^{g\delta_4}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (ii), then by Proposition 4.12, x_{r+5}^i exists and ${}_n\Phi_0^m(x_{r+5}^i) = v_{n-1}$, and so, $\tilde{\alpha}(x_{r+5}^i) = \tilde{s}_1^{g\delta_4}$ or $\tilde{\alpha}(x_{r+5}^i) = \tilde{s}_3^{g\delta_4}$. Then, $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_4}$, and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_1^{g\delta_4}$ or $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_3^{g\delta_4}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.v: $g \in G_3^{\{4\}} \cup G_3^{\{2,4\}}$.

From Case 4 of the previous claim,

i) $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_4^{g\delta_0}, \tilde{s}_3^{g\delta_0}, \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_0 \in F_2$ and $g\delta_5 \in F_0$ or

ii) $\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+5}^i \rangle) = \langle \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0}, \tilde{s}_0, \tilde{s}_1^{g\delta_3}, \tilde{s}_0, \tilde{s}_1^{g\delta_5} \rangle$ for some $g\delta_0 \in F_1$, $g\delta_3 \in F_1 \cup F_2$, and $g\delta_5 \in F_0$.

Case 2.b.v.1: $x = x_{r+4}^i$.

Then, $y = x_{r+5}^i$.

If (i) or (ii), then $\hat{\alpha}(x) = \hat{\alpha}(x_{r+4}^i) = \hat{s}_2^{g\delta_5}$ and $\hat{\alpha}(y) = \hat{\alpha}(x_{r+5}^i) = \hat{s}_{1+2}^{g\delta_5} = \hat{s}_3^{g\delta_5}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 2.b.v.2: $x = x_r^i$.

Then, $y = x_{r-1}^i$.

If (i), then by Proposition 4.12, x_{r-1}^i exists and ${}_n\Phi_0^m(x_{r-1}^i) = v_{n-1}$, and so, $\tilde{\alpha}(x_{r-1}^i) = \tilde{s}_3^{g\delta_0}$ or $\tilde{\alpha}(x_{r-1}^i) = \tilde{s}_5^{g\delta_0}$. Then, $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_0}$, and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_{3-2}^{g\delta_0} = \hat{s}_1^{g\delta_0}$ or $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_{5-2}^{g\delta_0} = \hat{s}_3^{g\delta_0}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

If (ii), then by Proposition 4.12, x_{r-1}^i exists and ${}_n\Phi_0^m(x_{r-1}^i) = v_{n-1}$, and so, $\tilde{\alpha}(x_{r-1}^i) = \tilde{s}_1^{g\delta_0}$ or $\tilde{\alpha}(x_{r-1}^i) = \tilde{s}_3^{g\delta_0}$. Then, $\hat{\alpha}(x) = \hat{\alpha}(x_r^i) = \hat{s}_2^{g\delta_0}$, and $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_1^{g\delta_0}$ or $\hat{\alpha}(y) = \hat{\alpha}(x_{r-1}^i) = \hat{s}_3^{g\delta_0}$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

Case 3: $x, y \in V({}_nX_m \setminus \bigcup G')$.

Case 3.a: $\tilde{\alpha}(x), \tilde{\alpha}(y) \in \tilde{S}^\delta \setminus \{\tilde{s}_0\}$ for some $\delta \in \{1, \dots, \tilde{n}_0\}$.

Since $\tilde{\alpha}$ is simplicial, then $\tilde{\alpha}(x)$ and $\tilde{\alpha}(y)$ are adjacent by Proposition 4.12, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent by definition of $\hat{\alpha}$.

Case 3.b: $\tilde{\alpha}(x) = \tilde{s}_0$.

Case 3.b.i: $\tilde{\alpha}(y) = \tilde{s}_1^\delta$ for some $\delta \in F_0$.

Then, ${}_n\Phi_0^m(y) = \tilde{\beta}(\tilde{\alpha}(y)) = \tilde{\beta}(\tilde{s}_1^\delta) \neq v_{n-1}$, and so by Proposition 4.12, there exist $i \in \{1, \dots, n\}$ and $z \in V({}_nX_m)$ so that z is adjacent to x and $z \neq y$ and $\langle z, x, y \rangle \subseteq {}_nA_m^i$. By Proposition 4.16, ${}_n\Phi_0^m(z) = v_{n-1}$, and so by Proposition 4.12, $z \in g$ for some $g \in G$.

Suppose $g \notin G_0$. Then, $z = x_{r+1}^i$ or $z = x_{r+3}^i$ where $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle$. Since ${}_n\Phi_0^m(y) \neq v_{n-1}$, then $\langle z, x, y \rangle = \langle x_{r+1}^i, x_r^i, x_{r-1}^i \rangle$ or $\langle z, x, y \rangle = \langle x_{r+3}^i, x_{r+4}^i, x_{r+5}^i \rangle$, and so, $\tilde{\alpha}(x_r^i) = \tilde{s}_0$ or $\tilde{\alpha}(x_{r+4}^i) = \tilde{s}_0$. Thus, $g \in G'$, contradicting $x \in g$.

Suppose $g \in G_0$. Then, $z = x_1^i$ and $x = x_2^i$ where $g = \langle x_0^i, x_1^i, x_2^i \rangle$, and so $g \in G'_0 \cup G''_0 \subseteq G'$, contradicting $x \in g$.

Thus, Case 3.b.i does not occur.

Case 3.b.ii: $\tilde{\alpha}(y) = \tilde{s}_1^\delta$ for some $\delta \in F_2$.

From Case 6 of the previous claim, $y \in g$ for some $g \in G'$, giving a contradiction.

Thus, Case 3.b.ii does not occur.

Case 3.b.iii: $\tilde{\alpha}(y) = \tilde{s}_1^\delta$ for some $\delta \in F_1$.

Then, $\hat{\alpha}(x) = \hat{s}_0$ and $\hat{\alpha}(y) = \hat{s}_1^\delta$, and so, $\hat{\alpha}(x)$ and $\hat{\alpha}(y)$ are adjacent.

CLAIM: $\hat{\beta}(\hat{s}_0) = v_0$.

By definition of $\hat{\beta}$, $\hat{\beta}(\hat{s}_0) = \tilde{\beta}(\tilde{s}_0)$, and by hypothesis, $\tilde{\beta}(\tilde{s}_0) = v_0$.

CLAIM: $\hat{\beta} \circ \hat{\alpha} = {}_n\Phi_0^m$.

Let $x \in V({}_nX_m)$.

Case 1: $x \in g$ for some $g \in G'$.

Case 1.a: $g \in G'_0 \cup G''_0$.

Let $g = \langle x_0^i, x_1^i, x_2^i \rangle$ for some $i \in \{1, \dots, n\}$. By Proposition 4.12, ${}_n\Phi_0^m(\langle x_0^i, x_1^i, x_2^i \rangle) = \langle v_0, v_{n-1}, v_0 \rangle$.

Case1.a.i: $g \in G'_0$.

From Case 2.a.i of the second claim, (i) or (ii) holds.

If (i), then $\overline{{}_n\Phi_0^m}(\langle x_0^i, x_1^i, x_2^i \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle)) = \overline{\beta}(\langle \tilde{s}_0, \tilde{s}_1^{g\delta_3}, \tilde{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\langle \hat{s}_0, \hat{s}_1^{g\delta_3}, \hat{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle))$.

If (ii), then $\overline{{}_n\Phi_0^m}(\langle x_0^i, x_1^i, x_2^i \rangle) = \langle v_0, v_{n-1}, v_0 \rangle = \overline{\beta}(\langle \hat{s}_0, \hat{s}_1^{g\delta_3}, \hat{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle))$.

Case1.a.ii: $g \in G''_0$.

From Case 2.a.ii of the second claim, (i),(ii), or (iii) holds.

If (i), then $\overline{{}_n\Phi_0^m}(\langle x_0^i, x_1^i, x_2^i \rangle) = \langle v_0, v_{n-1}, v_0 \rangle = \overline{\beta}(\langle \hat{s}_0, \hat{s}_1^{g\delta_3}, \hat{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle))$.

If (ii), then $\overline{{}_n\Phi_0^m}(\langle x_0^i, x_1^i, x_2^i \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle)) = \overline{\beta}(\langle \tilde{s}_0, \tilde{s}_1^{g\delta_3}, \tilde{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\langle \hat{s}_0, \hat{s}_1^{g\delta_3}, \hat{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle))$.

If (iii), then $\overline{{}_n\Phi_0^m}(\langle x_0^i, x_1^i, x_2^i \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle)) = \overline{\beta}(\langle \tilde{s}_2^{g\delta_3}, \tilde{s}_3^{g\delta_3}, \tilde{s}_4^{g\delta_3} \rangle) = \overline{\beta}(\langle \hat{s}_0, \hat{s}_1^{g\delta_3}, \hat{s}_2^{g\delta_3} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_0^i, x_1^i, x_2^i \rangle))$.

Case 1.b: $g \in \bigcup_{i=1}^4 G_i$.

Let $g = \langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle$ for some $i \in \{1, \dots, n\}$ and $r \in \{0, \dots, \ell_i\}$.

Case 1.b.i: $g \in G_1^{\{0\}} \cup G_1^{\{0,2\}}$.

From Case 2.b.i of the second claim, (i),(ii), or (iii) holds.

If (i), then $\overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle) \vee \overline{{}_n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1} \rangle \vee \overline{{}_n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \overline{\beta}(\langle \hat{s}_2^{g\delta_{-1}}, \hat{s}_1^{g\delta_{-1}}, \hat{s}_0, \hat{s}_1^{g\delta_4} \rangle) \vee \overline{\beta}(\overline{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \overline{\beta}(\langle \tilde{s}_3^{g\delta_4}, \tilde{s}_4^{g\delta_4} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \overline{\beta}(\langle \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \overline{\beta}(\overline{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, and if (ii) or (iii), then $\overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle) \vee \overline{{}_n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1} \rangle \vee \overline{{}_n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \overline{\beta}(\langle \hat{s}_2^{g\delta_{-1}}, \hat{s}_1^{g\delta_{-1}}, \hat{s}_0, \hat{s}_1^{g\delta_4} \rangle) \vee \overline{\beta}(\overline{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \overline{\beta}(\langle \tilde{s}_1^{g\delta_4}, \tilde{s}_2^{g\delta_4} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \overline{\beta}(\langle \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle) = \overline{\beta}(\overline{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \overline{\beta}(\overline{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) =$

$\bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, since $\hat{\beta}(\hat{s}_1^{g\delta_4}) = \tilde{\beta}(\tilde{s}_{1+2}^{g\delta_4}) = \tilde{\beta}(\tilde{s}_3^{g\delta_4}) = v_{n-1}$ if $g\delta_4 \in F_2$ and $\hat{\beta}(\hat{s}_1^{g\delta_4}) = \tilde{\beta}(\tilde{s}_1^{g\delta_4}) = v_{n-1}$ if $g\delta_4 \in F_1$.

Case 1.b.ii: $g \in G_1^{\{4\}} \cup G_1^{\{2,4\}}$.

From Case 2.b.ii of the second claim, (i), (ii), or (iii) holds.

If (i), then $\overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \overline{n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle) = \overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \langle v_{n-1}, v_0, v_{n-1}, v_0 \rangle = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i \rangle)) \vee \bar{\beta}(\langle \hat{s}_1^{g\delta_0}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle) = \bar{\beta}(\langle \hat{s}_4^{g\delta_0}, \hat{s}_3^{g\delta_0} \rangle) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \bar{\beta}(\langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0} \rangle) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, and if (ii) or (iii), then $\overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \overline{n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle) = \overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \langle v_{n-1}, v_0, v_{n-1}, v_0 \rangle = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i \rangle)) \vee \bar{\beta}(\langle \hat{s}_1^{g\delta_0}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle) = \bar{\beta}(\langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0} \rangle) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \bar{\beta}(\langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0} \rangle) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, since $\hat{\beta}(\hat{s}_1^{g\delta_0}) = \tilde{\beta}(\tilde{s}_{1+2}^{g\delta_0}) = \tilde{\beta}(\tilde{s}_3^{g\delta_0}) = v_{n-1}$ if $g\delta_0 \in F_2$ and $\hat{\beta}(\hat{s}_1^{g\delta_0}) = \tilde{\beta}(\tilde{s}_1^{g\delta_0}) = v_{n-1}$ if $g\delta_0 \in F_1$.

Case 1.b.iii: $g \in G_1^{\{0,4\}} \cup G_1^{\{0,2,4\}}$.

From Case 2.b.iii of the second claim, (i) or (ii) holds.

If (i) or (ii), then $\overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1}, v_0 \rangle = \bar{\beta}(\langle \hat{s}_2^{g\delta-1}, \hat{s}_1^{g\delta-1}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$.

Case 1.b.iv: $g \in G_2^{\{0\}} \cup G_2^{\{0,2\}}$.

From Case 2.b.iv of the second claim, (i) or (ii) holds.

If (i), then $\overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle) \vee \overline{n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1} \rangle \vee \overline{n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \bar{\beta}(\langle \hat{s}_2^{g\delta-1}, \hat{s}_1^{g\delta-1}, \hat{s}_0, \hat{s}_1^{g\delta_4} \rangle) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \bar{\beta}(\langle \hat{s}_3^{g\delta_4}, \hat{s}_4^{g\delta_4} \rangle) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \bar{\beta}(\langle \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, and if (ii), then $\overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle) \vee \overline{n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \langle v_0, v_{n-1}, v_0, v_{n-1} \rangle \vee \overline{n\Phi_0^m}(\langle x_{r+3}^i, x_{r+4}^i \rangle) = \bar{\beta}(\langle \hat{s}_2^{g\delta-1}, \hat{s}_1^{g\delta-1}, \hat{s}_0, \hat{s}_1^{g\delta_4} \rangle) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \bar{\beta}(\langle \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \bar{\beta}(\langle \hat{s}_1^{g\delta_4}, \hat{s}_2^{g\delta_4} \rangle) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+3}^i \rangle)) \vee \bar{\beta}(\bar{\alpha}(\langle x_{r+3}^i, x_{r+4}^i \rangle)) = \bar{\beta}(\bar{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, since $\hat{\beta}(\hat{s}_1^{g\delta_4}) = \tilde{\beta}(\tilde{s}_{1+2}^{g\delta_4}) = \tilde{\beta}(\tilde{s}_3^{g\delta_4}) = v_{n-1}$ if $g\delta_4 \in F_2$ and

$$\hat{\beta}(\hat{s}_1^{g\delta_4}) = \tilde{\beta}(\tilde{s}_1^{g\delta_4}) = v_{n-1} \text{ if } g\delta_4 \in F_1.$$

$$\text{Case 1.b.v: } g \in G_3^{\{4\}} \cup G_3^{\{2,4\}}.$$

From Case 2.b.v of the second claim, (i) or (ii) holds.

If (i), then $\overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \overline{{}_n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle) = \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \langle v_{n-1}, v_0, v_{n-1}, v_0 \rangle = \tilde{\beta}(\tilde{\alpha}(\langle x_r^i, x_{r+1}^i \rangle)) \vee \tilde{\beta}(\langle \hat{s}_1^{g\delta_0}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle) = \tilde{\beta}(\langle \tilde{s}_4^{g\delta_0}, \tilde{s}_3^{g\delta_0} \rangle) \vee \tilde{\beta}(\tilde{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \tilde{\beta}(\langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0} \rangle) \vee \tilde{\beta}(\tilde{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \tilde{\beta}(\tilde{\alpha}(\langle x_r^i, x_{r+1}^i \rangle)) \vee \tilde{\beta}(\tilde{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \tilde{\beta}(\tilde{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, and if (ii), then $\overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle) = \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \overline{{}_n\Phi_0^m}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle) = \overline{{}_n\Phi_0^m}(\langle x_r^i, x_{r+1}^i \rangle) \vee \langle v_{n-1}, v_0, v_{n-1}, v_0 \rangle = \tilde{\beta}(\tilde{\alpha}(\langle x_r^i, x_{r+1}^i \rangle)) \vee \tilde{\beta}(\langle \hat{s}_1^{g\delta_0}, \hat{s}_0, \hat{s}_1^{g\delta_5}, \hat{s}_2^{g\delta_5} \rangle) = \tilde{\beta}(\langle \tilde{s}_2^{g\delta_0}, \tilde{s}_1^{g\delta_0} \rangle) \vee \tilde{\beta}(\tilde{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \tilde{\beta}(\langle \hat{s}_2^{g\delta_0}, \hat{s}_1^{g\delta_0} \rangle) \vee \tilde{\beta}(\tilde{\alpha}(\langle x_{r+1}^i, x_{r+2}^i, \dots, x_{r+4}^i \rangle)) = \tilde{\beta}(\tilde{\alpha}(\langle x_r^i, x_{r+1}^i, \dots, x_{r+4}^i \rangle))$, since $\hat{\beta}(\hat{s}_1^{g\delta_0}) = \tilde{\beta}(\tilde{s}_{1+2}^{g\delta_0}) = \tilde{\beta}(\tilde{s}_3^{g\delta_0}) = v_{n-1}$ if $g\delta_0 \in F_2$ and $\hat{\beta}(\hat{s}_1^{g\delta_0}) = \tilde{\beta}(\tilde{s}_1^{g\delta_0}) = v_{n-1}$ if $g\delta_0 \in F_1$.

Case 2: $x \in V({}_nX_m \setminus \bigcup G')$.

Case 2.a: $\tilde{\alpha}(x) = \tilde{s}_0$.

$$\text{Then, } {}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_0) = \hat{\beta}(\hat{s}_0) = \hat{\beta}(\hat{\alpha}(x)).$$

Case 2.b: $\tilde{\alpha}(x) = \tilde{s}_k^\delta$ for some $\delta \in F_0$.

$$\text{Then, } {}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_k^\delta) = \hat{\beta}(\hat{s}_{k+2}^\delta) = \hat{\beta}(\hat{\alpha}(x)).$$

Case 2.c: $\tilde{\alpha}(x) = \tilde{s}_k^\delta$ for some $\delta \in F_2$.

Case 2.c.i: $k = 2$.

$$\text{Then, } {}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_2^\delta) = v_0 = \tilde{\beta}(\tilde{s}_0) = \hat{\beta}(\hat{s}_0) = \hat{\beta}(\hat{\alpha}(x)).$$

Case 2.c.ii: $k \in \{3, \dots, \tilde{k}_\delta\}$.

$$\text{Then, } {}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_k^\delta) = \hat{\beta}(\hat{s}_{k-2}^\delta) = \hat{\beta}(\hat{\alpha}(x)).$$

Case 2.d: $\tilde{\alpha}(x) = \tilde{s}_k^\delta$ for some $\delta \in F_1$.

$$\text{Then, } {}_n\Phi_0^m(x) = \tilde{\beta}(\tilde{\alpha}(x)) = \tilde{\beta}(\tilde{s}_k^\delta) = \hat{\beta}(\hat{s}_k^\delta) = \hat{\beta}(\hat{\alpha}(x)). \quad \square$$




Figure 4.5: (top) Factoring of ${}_5\Phi_0^m$ given by Theorem 3.3. (bottom) Factoring given by Proposition 4.17.

For Propositions 4.18 and 4.19 let $\varprojlim \{ {}_n Y_i, {}_n \Gamma_i^j \}_{i=0, j \geq i}^\infty$ be generated by $d[{}_n \phi] \circ {}_n \lambda, {}_n \phi, {}_n \phi, \dots$ (Definition 4.10).

Proposition 4.18. *Let $m \in \{1, 2, \dots\}$, $\tilde{n} \in \{1, \dots, n-1\}$, $\tilde{T}$ be a simple- $\tilde{n}$ -od, and $\tilde{\alpha} : {}_n Y_m \longrightarrow \tilde{T}$ and $\tilde{\beta} : \tilde{T} \longrightarrow {}_n Y_1$ be simplicial maps so that $\tilde{\beta} \circ \tilde{\alpha} = {}_n \Gamma_1^m$ and $\tilde{\beta}(\tilde{s}_0) = v_0$ where $\tilde{s}_0$ is the branch point of $\tilde{T}$. Then, there exist a simple- $\tilde{n}$ -od T and simplicial maps $\alpha : {}_n X_{m-1} \longrightarrow T$ and $\beta : T \longrightarrow {}_n X_0$ so that $\beta \circ \alpha = {}_n \Phi_0^{m-1}$ and $\beta(s_0) = v_0$ where s_0 is the branch point of T .*

Proof. Let $\tilde{S}^1, \tilde{S}^2, \dots, \tilde{S}^{\tilde{n}}$ be $\tilde{n}$ arcs so that $\bigcup_{i=1}^{\tilde{n}} \tilde{S}^i = \tilde{T}$ and $\tilde{S}^i \cap \tilde{S}^j = \{\tilde{s}_0\}$ for each distinct $i, j \in \{1, \dots, \tilde{n}\}$, and so $\tilde{s}_0$ is an endpoint of $\tilde{S}^i$ for each $i \in \{1, \dots, \tilde{n}\}$.

CLAIM: $\tilde{\alpha}^{-1}(\tilde{S}^i)$ is a subdivision of a subgraph ${}_n X_{m-1}^i$ of ${}_n X_{m-1}$ for each $i \in \{1, \dots, \tilde{n}\}$.

Let $i \in \{1, \dots, \tilde{n}\}$ and $x, y \in V({}_n X_{m-1})$ so that x and y are distinct and adjacent and $(\langle x, y \rangle, {}_n Y_m) \cap \tilde{\alpha}^{-1}(\tilde{S}^i) \neq \emptyset$. Since $\tilde{\beta}(\tilde{s}_0) = v_0$ and ${}_n \Gamma_1^m$ is a subdivision of ${}_n \Phi_0^{m-1}$ matching ${}_n Y_1$, then $\tilde{\alpha}^{-1}(\tilde{s}_0) \subseteq V({}_n X_{m-1})$. Suppose $\tilde{\alpha}(x) \in \tilde{S}^i$. If $\tilde{\alpha}(y) \notin \tilde{S}^i$, then $(\langle x, y \rangle, {}_n Y_m) \cap \tilde{\alpha}^{-1}(\tilde{S}^i) = \{x\}$, and if $\tilde{\alpha}(y) \in \tilde{S}^i$, then $(\langle x, y \rangle, {}_n Y_m) \subseteq \tilde{\alpha}^{-1}(\tilde{S}^i)$. Suppose $\tilde{\alpha}(x) \notin \tilde{S}^i$. If $\tilde{\alpha}(y) \notin \tilde{S}^i$, then $(\langle x, y \rangle, {}_n Y_m) \cap \tilde{\alpha}^{-1}(\tilde{S}^i) = \emptyset$, and if $\tilde{\alpha}(y) \in \tilde{S}^i$, then $(\langle x, y \rangle, {}_n Y_m) \cap \tilde{\alpha}^{-1}(\tilde{S}^i) = \{y\}$.

Thus, the claim holds.

By Proposition 5.13 in [14], for each $i \in \{1, \dots, \tilde{n}\}$, there exist an arc S^i , simplicial maps $\beta^i : S^i \longrightarrow {}_n \Phi_0^{m-1}({}_n X_{m-1}^i)$ and $\alpha^i : {}_n X_{m-1}^i \longrightarrow S^i$ so that $\beta^i \circ \alpha^i = {}_n \Phi_0^{m-1}|_{{}_n X_{m-1}^i}$, and an endpoint s_0^i of S^i so that $\beta^i(s_0^i) = \tilde{\beta}(\tilde{s}_0)$ and $\alpha^{i-1}(s_0^i) = \tilde{\alpha}^{-1}(\tilde{s}_0)$.

Let $T = \bigcup_{i=1}^{\tilde{n}} S^i / \sim$, where $\sim$ identifies s_0^i to s_0 for all $i \in \{1, \dots, \tilde{n}\}$, and $q : \bigcup_{i=1}^{\tilde{n}} S^i \longrightarrow \bigcup_{i=1}^{\tilde{n}} S^i / \sim$ be the quotient map, and define $\beta : T \longrightarrow {}_n X_0$ as $\beta(y) = \beta^i(x)$ if $y = q(x)$ for some $x \in V(S^i \setminus \{s_0^i\})$ and $i \in \{1, \dots, \tilde{n}\}$ and $\beta(s_0) = \tilde{\beta}(\tilde{s}_0)$, and $\alpha : {}_n X_{m-1} \longrightarrow T$ as $\alpha(x) = q(\alpha^i(x))$ if $x \in V({}_n X_{m-1}^i)$.

Since β^i is simplicial and $\beta^i(s_0^i) = \tilde{\beta}(\tilde{s}_0)$ for all $i \in \{1, \dots, \tilde{n}\}$, then β is simplicial.

CLAIM: α is well-defined.

Let $x \in V({}_nX_{m-1}^i \cap {}_nX_{m-1}^j)$ for distinct $i, j \in \{1, \dots, \tilde{n}\}$. Then, $x \in \tilde{\alpha}^{-1}(\tilde{s}_0)$, and so, $q(\alpha^i(x)) = q(s_0^i) = s_0 = q(s_0^j) = q(\alpha^j(x))$.

CLAIM: α is simplicial.

Let $x, y \in V({}_nX_{m-1})$ be distinct and adjacent.

By the first claim, $x, y \in {}_nX_{m-1}^i$ for some $i \in \{1, \dots, \tilde{n}\}$. Since q is simplicial, α^i is simplicial, and ${}_n\Phi_0^{m-1}$ is light, then $\alpha(x) = q(\alpha^i(x))$ and $\alpha(y) = q(\alpha^i(y))$ are adjacent.

CLAIM: $\beta \circ \alpha = {}_n\Phi_0^{m-1}$.

Let $x \in V({}_nX_{m-1})$. Then, $x \in V({}_nX_{m-1}^i)$ for some $i \in \{1, \dots, \tilde{n}\}$. If $x \in \tilde{\alpha}^{-1}(\tilde{s}_0)$, then ${}_n\Phi_0^{m-1}(x) = \beta^i(\alpha^i(x)) = \beta^i(s_0^i) = \tilde{\beta}(\tilde{s}_0) = \beta(s_0) = \beta(q(s_0^i)) = \beta(q(\alpha^i(x))) = \beta(\alpha(x))$, and if $x \notin \tilde{\alpha}^{-1}(\tilde{s}_0)$, then $\alpha^i(x) \in V(S^i \setminus \{s_0^i\})$ and ${}_n\Phi_0^{m-1}(x) = \beta^i(\alpha^i(x)) = \beta(q(\alpha^i(x))) = \beta(\alpha(x))$.

CLAIM: $\beta(s_0) = v_0$.

By definition, $\beta(s_0) = \tilde{\beta}(\tilde{s}_0)$, and by hypothesis, $\tilde{\beta}(\tilde{s}_0) = v_0$. $\square$

Proposition 4.19. *There do not exist a simple- $\tilde{n}$ -od T and simplicial maps $\alpha : {}_nX_m \longrightarrow T$ and $\beta : T \longrightarrow {}_nX_0$ so that $\beta \circ \alpha = {}_n\Phi_0^m$ and $\beta(s_0) = v_0$, where s_0 is the branch point of T , for each $m \in \{1, 2, \dots\}$ and for each $\tilde{n} \in \{1, \dots, n-1\}$.*

Proof. Let $\tilde{n} \in \{1, \dots, n-1\}$ and ${}_nA_1^i = \langle x_0^i, x_1^i, \dots, x_{\ell_i}^i \rangle$ for some $\ell_i \in \{1, 2, \dots\}$ for each $i \in \{1, \dots, n\}$ where $x_0^i = v_0$ for all $i \in \{1, \dots, n\}$.

Let $m = 1$ and suppose there exist a simple- $\tilde{n}$ -od T and simplicial maps $\alpha : {}_nX_1 \longrightarrow T$ and $\beta : T \longrightarrow {}_nX_0$ so that $\beta \circ \alpha = {}_n\Phi_0^1$ and $\beta(s_0) = v_0$ where s_0 is the branch point of T .

Since $\overline{{}_n\Phi_0^1}({}_nA_1^i) = \langle v_0, v_{n-1}, v_0, v_{p^{i-1}(1)}, v_0, \underline{v_{p^{i-1}(2)}, v_0}, \dots, v_{p^{i-1}(n-2)}, v_0}, v_n, v_{n+2} \rangle$ for each $i \in \{1, \dots, n-3\}$, $\overline{{}_n\Phi_0^1}({}_nA_1^{n-2}) = \langle v_0, v_{n-1}, v_{n+1} \rangle$, $\overline{{}_n\Phi_0^1}({}_nA_1^{n-1}) = \langle v_0, v_{n-1}, v_0, v_n, v_{n+2} \rangle$, and $\overline{{}_n\Phi_0^1}({}_nA_1^n) = \langle v_0, v_{n-1}, v_0, v_{n-2}, v_0, v_1, v_0, \underline{v_2, v_0}, \dots, \underline{v_{n-3}, v_0}, v_n, v_{n+2} \rangle$, then ${}_n\Phi_0^1(x_{t_i}^i) = {}_n\Phi_0^1(x_{t_j}^j)$ for some $t < t_j$ where $x_{t_i}^i, x_{t_j}^j \in \{x_2^{n-2}, x_3^k : k \in \{1, \dots, n\} \setminus \{n-2\}\}$ are distinct, giving a contradiction.

Let $m > 1$, and suppose the claim holds for all $\tilde{m} \in \{1, \dots, m-1\}$. Suppose


there exist a simple- $\tilde{n}$ -od T and simplicial maps $\alpha : {}_nX_m \longrightarrow T$ and $\beta : T \longrightarrow {}_nX_0$ so that $\beta \circ \alpha = {}_n\Phi_0^m$ and $\beta(s_0) = v_0$ where s_0 is the branch point of T .

Since ${}_nX_m$ is connected, α may be assumed to be surjective. By Proposition 4.17, there exist $\tilde{n}_1, \hat{T}, \hat{\alpha}$, and $\hat{\beta}$ as defined in Proposition 4.17. Since $\hat{\beta}(\bigcup_{i=1}^{\tilde{n}_1} \langle \hat{s}_0, \hat{s}_1^i \rangle) = \langle v_0, v_{n-1} \rangle$, then $\bigcup_{i=1}^{\tilde{n}_1} \langle \hat{s}_0, \hat{s}_1^i \rangle \subseteq c_0^*$ and $d[\hat{\beta}](c_0) = a_{n-1}$ where c_0 is a vertex of $D(\hat{\beta}, \hat{T})$. So, if $c_i, c_j \in V(D(\hat{\beta}, \hat{T}))$ are adjacent, then $c_i^* \cup c_j^* \subseteq \hat{S}^\delta$ for some $\delta \in \{1, \dots, \tilde{n}\}$, or $c_k = c_0$ for some $k \in \{i, j\}$ and $c_\ell^* \subseteq \hat{S}^\delta$ for some $\delta \in \{1, \dots, \tilde{n}\}$ where $\ell \in \{i, j\}$ and $\ell \neq k$, giving $D(\hat{\beta}, \hat{T})$ is a simple- $\tilde{n}$ -od with branch point c_0 .

Since $\hat{\alpha}$ is surjective by definition, $d[\hat{\beta}, \hat{\alpha}]$ is surjective. By Proposition 4.7, Proposition 2.6 in [14], and Theorem 4.3 in [14], there exists a simplicial map $\sigma : D(\hat{\beta}, \hat{T}) \longrightarrow D({}_n\Phi_0^1, {}_nX_1)$ so that $\sigma \circ d[\hat{\beta}, \hat{\alpha}] = d[{}_n\Phi_0^1, {}_n\Phi_1^m]$. By the proof of Proposition 4.7, $d[{}_n\phi]^{-1}(a_{n-1}) = \{b_0\}$, and so, since $d[\hat{\beta}](c_0) = a_{n-1}$, $\sigma(c_0) = b_0$.

By Proposition 4.6, Proposition 4.9, and Theorem 5.11 in [14], $\varprojlim \{D({}_n\Phi_0^i, {}_nX_i), d[{}_n\Phi_0^i, {}_n\Phi_i^j]\}_{i=0, j \geq i}^\infty$ is isomorphic to $\varprojlim \{{}_nY_i, {}_n\Gamma_i^j\}_{i=0, j \geq i}^\infty$ (Definition 4.11). Thus, there exist a simple- $\tilde{n}$ -od $\tilde{T}$ and simplicial maps $\tilde{\alpha} : {}_nY_m \longrightarrow \tilde{T}$ and $\tilde{\beta} : \tilde{T} \longrightarrow {}_nY_1$ so that $\tilde{\beta} \circ \tilde{\alpha} = {}_n\Gamma_1^m$ and $\tilde{\beta}(\tilde{s}_0) = v_0$ where $\tilde{s}_0$ is the branch point of $\tilde{T}$.

By Proposition 4.18, there exist a simple- $\tilde{n}$ -od T and simplicial maps $\alpha : {}_nX_{m-1} \longrightarrow T$ and $\beta : T \longrightarrow {}_nX_0$ so that $\beta \circ \alpha = {}_n\Phi_0^{m-1}$ and $\beta(s_0) = v_0$ where s_0 is the branch point of T , contradicting the induction hypothesis. Therefore, the claim follows by induction. $\square$


 Figure 4.6: Factoring of ${}_5\Gamma_1^m$

Proposition 4.20. ${}_nK$ is not simple- $(n-1)$ -od-like.

Proof. If ${}_nK$ is simple- $(n-1)$ -od-like, then the conclusion of Theorem 3.3 for $j = 0$ contradicts Proposition 4.19. $\square$

Proposition 4.21. ${}_nK$ is indecomposable.

Proof. The claim follows from Proposition 4.3, Proposition 4.4, Proposition 4.20, and Corollary 1 in [3]. $\square$

Theorem 4.1. For each integer n greater than or equal to 3, there exists a simple- n -od-like continuum $({}_nK)$ having the properties of not being simple- $(n-1)$ -od-like and of every proper nondegenerate subcontinuum being an arc.

Proof. The claim follows from Propositions 4.3 and 4.20. $\square$

CHAPTER V

FURTHER QUESTIONS

This dissertation provides, for each integer $n \geq 3$, a simple- n -od-like continuum ${}_nK$ which is not simple- $(n-1)$ -od-like and whose every proper nondegenerate subcontinuum is an arc. Examples for the case $n = 5$ and higher were not proven previously. The existence of such continua is related to the problem of being able to distinguish among tree-like continua those that are chainable. A fundamental open question in this area is the following (Question 5.1) due to L. Mohler. One partial positive answer has been given by L.G. Oversteegen in [17] for continua satisfying the additional conditions of being the continuous image of a chainable continuum under an induced map and of having surjective semispan equal to zero. Another partial positive answer has been given by P. Minc in [16], as another application of the combinatorial machinery developed in [14], for continua which are the inverse limits of trees with simplicial bonding maps.

Question 5.1 (Problem 16 [12]). *Is every atriodic tree-like continuum that is the continuous image of a chainable continuum chainable?*

Examining possible patterns for nested intersections of covers as determined by the pattern of the bonding map ${}_n\phi$ (for $n \geq 4$), the examples presented here do not appear to be planar. The question of just how “simple” examples with these properties could be, corresponding to questions regarding the complexity of subcontinua of the plane, is raised in the following.

Question 5.2. *For each integer n greater than 3, does there exist a simple- n -od-like planar continuum having the properties of not being simple- $(n-1)$ -od-like and of every proper nondegenerate subcontinuum being an arc (being atriodic)?*

Do there exist examples with the above-mentioned properties, replacing arc with pseudo-arc? The combinatorial techniques employed here, being dependent on a

simplicial setting, could not be directly utilized in showing that such a given example is not simpler (Definition 1.16). The case for $n = 3$ is known, as mentioned in the Introduction, with an example in [8], constructed so as to be continuously mapped onto the example in [7]. This implies nonchainability of the example in [8] since the example in [7] is not the continuous image of a chainable continuum. This does not naturally generalize to a method for showing that a hereditarily indecomposable simple- n -od-like continuum is not simple- $(n-1)$ -od-like. Specifically, can the bonding map for ${}_nK$ be modified, in a manner similar to that done to the bonding map in [7] to give the example in [8], to produce a continuum which is not simple- $(n-1)$ -od-like? What techniques could be used to recognize that such a continuum is not simple- $(n-1)$ -od-like?

Question 5.3. *For each integer n greater than 3, does there exist a simple- n -od-like (planar) continuum having the properties of not being simple- $(n-1)$ -od-like and of being hereditarily indecomposable (every proper nondegenerate subcontinuum being a pseudo-arc)?*

The following question concerns the existence of more “complicated” atriodic examples known to be like a certain graph and not like any simpler graph (Definition 1.15). For trees, in constructing an atriodic continuum like a given tree and not like anything simpler, the examples ${}_nK$ could be a way to control the order of the branching.

Question 5.4. *For a graph (tree) G , does there exist a G -like continuum having the properties of not being H -like, for all graphs H simpler (Definition 1.15) than G , and of being atriodic (being hereditarily indecomposable)?*

Question 5.5. *Does there exist a continuum K , with the property of being atriodic (of every proper nondegenerate subcontinuum being an arc), which is not simple- n -od-like for each positive integer n and which for each $\epsilon > 0$ there exist a positive integer m and an open cover $\mathcal{U}$ of K so that $\text{mesh}(\mathcal{U}) < \epsilon$ and the nerve of $\mathcal{U}$ is a simple- m -od?*

If v in a continuum K is a branch point of K of order n , then v is a branch point of K of order $n + 1$. If K is simple- $(n - 1)$ -od-like, is v a branch point of K of order $n - 1$? For ${}_nK$, by construction, $(v_0, v_0, v_0, \dots)$ is a branch point of ${}_nK$ of order n . In supposing ${}_nK$ is simple- $(n - 1)$ -od-like, by Proposition 3.2, $(v_0, v_0, v_0, \dots)$ is “close” to being a branch point of ${}_nK$ of order $n - 1$. Although sufficient in showing ${}_nK$ not being simple- $(n - 1)$ -od-like, the argument could be made more concise in the case of $(v_0, v_0, v_0, \dots)$ necessarily being a branch point of ${}_nK$ of order $n - 1$.

Question 5.6. *Does there exist a continuum K with $v \in K$ so that, for some integer n , v is a branch point of K of order n , v is not a branch point of K of order $n - 1$, and K is simple- $(n - 1)$ -od-like?*

BIBLIOGRAPHY

- [1] Anderson, R.D. *Hereditarily indecomposable plane continua* (Abstract 236). Bull. Amer. Math. Soc. 57 (1951), 185.
- [2] Bing, R.H. *Concerning hereditarily indecomposable continua*. Pacific J. Math. 1 (1951), 43-51.
- [3] Bing, R.H. *Snake-like continua*. Duke Math. J. 18 (1951), 653-663.
- [4] Cook, H., Ingram, W.T., and Lelek, A. *Eleven annotated problems about continua*. Open Problems in Topology. North-Holland, Amsterdam, 1990, 295-302.
- [5] Davis, J.F and Ingram, W.T. *An atriodic tree-like continuum with positive span which admits a monotone mapping to a chainable continuum*. Fund. Math. 131 (1988), 13-24.
- [6] Ingram, W.T. *Decomposable circle-like continua*. Fund. Math. 63 (1968), 193-198.
- [7] Ingram, W.T. *An atriodic tree-like continuum with positive span*. Fund. Math. 77 (1972), 99-107.
- [8] Ingram, W.T. *Hereditarily indecomposable tree-like continua*. Fund. Math. 103 (1979), 61-64.
- [9] Kato, H. and Ye, X. *On Burgess's theorem and related problems*. Proc. Amer. Math. Soc. 128 (2000), 2501-2506.
- [10] Lelek, A. *Disjoint mappings and the span of spaces*. Fund. Math. 55 (1964), 199-214.
- [11] Lewis, W. *Periodic homeomorphisms of chainable continua*. Fund. Math. 117 (1983), 81-84.
- [12] Lewis, W. *Continuum theory problems*. Top. Proc. 8 (1983), 361-394.
- [13] Mayer, J.C. *Principal embeddings of atriodic plane continua*. Continua, Decompositions, Manifolds. University of Texas Press, 1983, 34-61.
- [14] Minc, P. *On simplicial maps and chainable continua*. Topology and its Applications 57 (1994), 1-21.
- [15] Minc, P. *An atriodic simple-4-od-like continuum which is not simple-triod-like*. Trans. Amer. Math. Soc. 338 (1993), 537-552.

- [16] Minc, P. *On weakly chainable inverse limits with simplicial bonding maps.* Proc. Amer. Math. Soc. 119 (1993), 281-289.
- [17] Oversteegen, L.G. *On span and chainability of continua.* Houston J. Math. 15 (1989), 573-593.
- [18] Young, S.W. *A note concerning a continuum of J.C. Mayer.* Continua, Decompositions, Manifolds. University of Texas Press, 1983, 83-84.
- [19] Young, S.W. *On the relative complexity of a tree-like continuum and its proper subcontinua.* Continua, Decompositions, Manifolds. University of Texas Press, 1983, 85-95.

APPENDIX

SOME BONDING MAPS




Figure A.1: The bonding map from [7]

The bonding map above is used by W.T. Ingram as the single bonding map in the construction of an inverse limit, published in 1972, being the first proven counterexample to a question of Bing from 1951 as to whether every atriodic nonseparating plane continuum is chainable. Nonchainability of Ingram's continuum follows from his proof of positive span of the continuum, and the nature of the bonding map ensures every proper nondegenerate subcontinuum being an arc, implying atriodicity. As evident by examining the pattern above, the continuum is embeddable in the plane. Ingram's continuum is the first continuum which, by construction, is simple-3-od-like and shown not to be simple-2-od-like.




Figure A.2: The bonding map from [5]

The bonding map above is used by J.F. Davis and W.T. Ingram as the single bonding map in the construction of an inverse limit, published in 1988, with the properties of having positive span and of having every proper nondegenerate subcontinuum as an arc in common with the previous example, with similar techniques utilized in demonstrating positive span. The example is constructed to satisfy an additional property. The nature of the bonding map allows for a continuous map to be induced from the Davis-Ingram continuum to a chainable continuum, having only one nondegenerate point inverse which is an arc. The Davis-Ingram continuum is the first known example, having the previous example's properties, admitting a continuous monotone map to a chainable continuum.




Figure A.3: The bonding map from [15]

The bonding map above is used by P. Minc as the single bonding map in the construction of an inverse limit, published in 1993, with the properties of not being simple-3-od-like and of being atriodic. As with the others, the nature of the bonding map ensures every proper nondegenerate subcontinuum being an arc. In showing the continuum to be not simple-3-od-like, alternate techniques to span are needed. In so doing, Minc adapts combinatorial techniques from [14]. Minc's continuum is the first continuum which, by construction, is simple-4-od-like and shown not to be simple-3-od-like.